


**Site Description Data Sheet**


Site Code: \_\_\_\_\_ Orientation: \_\_\_\_\_ Date: \_\_\_\_\_  
 Recorder: \_\_\_\_\_ Crew: \_\_\_\_\_  
 GPS error: \_\_\_\_\_ Location: \_\_\_\_\_ Plot Size: \_\_\_\_\_  
 Slope: \_\_\_\_\_ ° Aspect: \_\_\_\_\_ ° Elevation: \_\_\_\_\_ m Terrain: E R G B  
 Meso Position: c u m l t d l Drainage: E G R Structure: SS MS C  
 Habitat F WL SW PP ST GT AT Moisture: vx x sx sm m sh h sH H

**Site Diagram:**



PC - percent cover  
 MVH - max veg height  
 CC - community composition (detailed)

**Location Map:**




**Berry Plots:**

	Cloudberry Rubcha	Crowberry Empnig	Blueberry Vaculi	Cranberry Vacvit
Overall Site				
Berry Plot 1				
Berry Plot 2				
Berry Plot 3				
Berry Plot 4				
Berry Plot 5				
Berry Plot 6				
Berry Plot 7				
Berry Plot 8				

Please indicate the overall abundance of each berry species above (dominant, sparse, not present, or not in season).  
 Mark an 'X' for each berry plot you establish under the species that it has been established for.

NW corner



Place

AK - Aklavik  
 IN - Inuvik  
 TK - Tuk

Site Code

Place - Habitat - Number

Trees

Sw=white spruce  
 Sb=black spruce  
 Bl=subalpine fir  
 Lt=tamarack  
 Pl=lodgepole pine  
 Pj=jack pine  
 At=trembling aspen  
 Ab=balsam poplar  
 Ep=paper birch  
 Ea=alaska paper birch

<u>Terrain</u>	<u>Structure</u>	<u>Habitat</u>
Even	single story	F=forest
Rolling		WL=woodland
Gullied	muti story	SW=sedge wetland
Broken	complex	PP=polygonal peatland
		ST=shrub tundra
<u>Drainage</u>		GT=graminoid tundra
Extreme		AT=alpine tundra
Good		
Restricted		

Very Xeric

water removed extremely rapidly in relation to supply  
 soil remains moist for a negligible time after precipitation  
 Primary water source is precipitation

Texture: very coarse; abundant coarse fragments

Xeric

water removed very rapidly in relation to supply  
 soil remains moist for a brief time after precipitation  
 primary water source is precipitation

Texture: coarse fragments

Subxeric

water removed rapidly in relation to supply  
 soil remains moist for a short period of time after precipitation  
 primary water source is precipitation

Texture: coarse to moderately coarse fragments

Submesic

water removed readily in relation to supply  
 water available for moderately short time periods after precipitation  
 primary water source is precipitation

Texture: moderately coarse

Mesic

water removed somewhat slowly in relation to supply  
 soil may remain moist for a significant, but sometimes short, time period  
 after ppt

primary water source is precipitation, however may be from limited  
 seepage in coarser texture soils

Texture: moderate to fine; few coarse fragments

Subhygric

water removed slowly enough to keep the soil wet for a significant part  
 of the growing season

some temporary seepage and possible mottling below 20 cm

primary water sources are precipitation and seepage

Texture: variable, depending on seepage

Hygric

water removed slowly enough to keep the soil wet for most of the  
 growing season

permanent seepage and mottling present; possibly weak gleying

primary water source is seepage

Texture: variable, depending on seepage

Subhydryc

water removed slowly enough to keep the water table at or near the  
 surface for most of the year

permanent seepage ≥ 30 cm from surface; gleyed soils

primary water sources are seepage or permanent water table

Texture: variable, depending on seepage

Hydryc

water table at or above the soil surface all year

gleyed / mottled soils

primary water source is the water table

