

# Environment

**Lithosphere** – solid ground – 100 km thick

**Hydrosphere** – ocean, rivers, lakes, ground water

**Atmosphere** – gases that surround us

**Table 14.1** The 10 Most Abundant Elements of the Earth's Crust

Element	Average Percentage (by weight)
Oxygen	46
Silicon	28
Aluminum	8
Iron	6
Calcium	4
Sodium	3
Magnesium	2
Potassium	2
Titanium	0.6
Hydrogen	0.1

**Table 14.2** The 10 Most Abundant Elements of the Earth's Oceans

Element	Average Percentage <sup>a</sup> (by weight)
Oxygen	86
Hydrogen	11
Chlorine	2
Sodium	1
Magnesium	0.1
Sulfur	0.09
Calcium	0.04
Potassium	0.04
Bromine	0.006
Carbon	0.003

<sup>a</sup>The total percentage is more than 100 because of the effects of rounding.

**Table 14.3** The 10 Most Abundant Gases of the Earth's Atmosphere

Gas	Average Percentage (by volume in dry air)
N <sub>2</sub>	78
O <sub>2</sub>	21
Ar	0.9
CO <sub>2</sub>	0.03
Ne	0.002
He	0.0005
CH <sub>4</sub>	0.0002
Kr	0.0001
H <sub>2</sub>	0.00005
N <sub>2</sub> O	0.00005

*The Extraordinary Chemistry of Ordinary Things, 4<sup>th</sup> Ed.*

# Chemistry in the Atmosphere

Smog




Los Angeles



Beijing

*wikipedia*


## Atmosphere – not all gases are equally distributed



wikipedia

Oxygen Content of Earth's Atmosphere  
During the Course of the Last Billion Years

Concentrations are **not stable** over *long* time: initially zero but cyanobacteria photosynthesis eventually produced **excess O<sub>2</sub>** then photosynthetic plants caused a second rapid increase




Limited by fires

wikipedia

Chemistry in Context 6<sup>th</sup> Edition, ACS, McGraw-Hill

# Tropospheric Pollutants: Ozone in the troposphere fluctuates during the day

**Ozone – secondary pollutant – formed with sun light**






**Figure 1.16**

Ozone level maps for a summer day in California, July 2006.

*The Extraordinary Chemistry of Ordinary Things, 4<sup>th</sup> Ed.*

# Pollutants from Coal

## Oxides of sulfur or nitrogen




*The Extraordinary Chemistry of Ordinary Things, 4<sup>th</sup> Ed.*

*Chemistry in Context 6<sup>th</sup> Edition, ACS, McGraw-Hill*


# Acid rain

**Sulfuric acid** –  $\text{H}_2\text{SO}_4$


**Nitric acid** –  $\text{HNO}_3$



Falling raindrop  
absorbs atmospheric  
carbon dioxide . . .



. . . which reacts with  
the water of the  
raindrop . . .



. . . to form  
carbonic acid.

1920's




1990's




*The Extraordinary Chemistry of Ordinary Things, 4<sup>th</sup> Ed.*

## Hydrogen ion concentration as pH from measurements made at the Central Analytical Laboratory, 1999



National Atmospheric Deposition Program/National Trends Network  
<http://nadp.sws.uiuc.edu>

## Hydrogen ion concentration as pH from measurements made at the Central Analytical Laboratory, 2013



### Sites not pictured:





Alaska 01	5.3
Alaska 02	5.2
Alaska 03	5.3
Alaska 97	5.1
Puerto Rico 20	5.2
British Columbia 22	4.4
Saskatchewan 21	5.5

National Atmospheric Deposition Program/National Trends Network  
<http://nadp.isws.illinois.edu>



# Air Pollution

## Photochemical smog




## Thermal inversion



*The Extraordinary Chemistry of Ordinary Things, 4<sup>th</sup> Ed.*

# World NO<sub>2</sub> distribution: 2009



## More pollutants

CO >>> odorless, tasteless and invisible

### Volatile organic compounds and volatile organic solvents

**Table 14.4 Major Sources of Air Pollution in the United States, 1998**

	Millions of Tons				Total
	Carbon Monoxide	Nitrogen Oxides	Sulfur Dioxide	Volatile Organic Compounds	
Fuel combustion					
Highway vehicles	50.4	7.8	0.3	5.3	63.8
Electric utilities	0.4	6.1	13.2	0.1	19.8
All other nonvehicle sources	5.0	4.1	3.5	0.8	13.4
Industrial emissions (other than fuel combustion)	3.6	0.8	1.5	1.4	7.3
Waste disposal and recycling	1.2	0.1	0	0.4	1.7

SOURCE: *U.S. Statistical Abstracts, 2000*, Table 394.

*The Extraordinary Chemistry of Ordinary Things, 4<sup>th</sup> Ed.*




See also: *Real-time World Air Quality Index*: <https://waqi.info/>


# SO<sub>x</sub> and NO<sub>x</sub>

Byproduct of energy production: combustion of hydrocarbons containing N and S impurities

## Electrostatic precipitator






## Filtration



*The Extraordinary Chemistry of Ordinary Things, 4<sup>th</sup> Ed.*

## Scrubbers




## The Stratospheric Ozone Layer: Earth's Protective Blanket

UV A: 315-400 nm

UV B: 280-315 nm

UV C: 200-280 nm



University Corporation for Atmospheric Research (UCAR): courtesy of NASA

# Chlorofluorocarbons: unintended consequences and a success story for action

Ozone destruction in the stratosphere – CFCs, •OH, •NO

# of F  
↓  
Freon 12  
↑  
# of H + 1  
(see later)

Table 2.6 Two Important Chlorofluorocarbons

Freon 11 (CFC-11)	Freon 12 (CFC-12)
$\text{CCl}_3\text{F}$ trichlorofluoromethane	$\text{CCl}_2\text{F}_2$ dichlorodifluoromethane
$\begin{array}{c} \text{:}\ddot{\text{F}}\text{:} \\   \\ \text{:}\ddot{\text{Cl}}\text{--C--}\ddot{\text{Cl}}\text{:} \\   \\ \text{:}\ddot{\text{Cl}}\text{:} \end{array}$	$\begin{array}{c} \text{:}\ddot{\text{F}}\text{:} \\   \\ \text{:}\ddot{\text{Cl}}\text{--C--}\ddot{\text{F}}\text{:} \\   \\ \text{:}\ddot{\text{Cl}}\text{:} \end{array}$


## Uses of CFC:

- refrigerant
- aerosol for spray cans
- gas to expand plastic foam
- solvent for oil and grease
- sterilizer for surgical instruments

Halons contain bromine (Br) or fluorine (F) to replace some of the Cl

*Chemistry in Context 6<sup>th</sup> Edition, ACS, McGraw-Hill*

## CFCs are destroyed by UV light at high altitudes




From: On-line Textbook, <http://biologytb.net23.net/text/index.html>

**At 20 -25 km  $\text{Cl}\bullet$  forms  $\text{HCl}$**


**Most  $\text{O}_3$  destruction occurs at 40 km**






Chemistry in Context 6<sup>th</sup> Edition, ACS, McGraw-Hill

## Milestones in the History of Stratospheric Ozone Depletion




EESC: equiv. effective stratospheric Cl

<http://www.theozonehole.com/>



## Banning OF CFC's: *An Ozone Recovery Story*

*2017 Antarctic ozone hole was the smallest since 1988*



University Corporation for Atmospheric Research (UCAR): courtesy of NASA

C. Bohne/D. Berg Copyright 2011-19 – For use in the Uvic Chem 400 course only – Spring 2019

## Replacement CFCs

- **Correct boiling point between -10 and -30 °C**
- **Not toxic**
- **Not too stable**
- **Not flammable**

Table 2.7

Two Important Hydrochlorofluorocarbons

HCFC-22	HCFC-141b
$\text{CHClF}_2$ chlorodifluoromethane	$\text{C}_2\text{H}_3\text{Cl}_2\text{F}$ dichlorofluoroethane
$  \begin{array}{c}  \text{:}\ddot{\text{F}}\text{:} \\    \\  \text{H}-\text{C}-\ddot{\text{F}}\text{:} \\    \\  \text{:}\ddot{\text{Cl}}\text{:}  \end{array}  $	$  \begin{array}{c}  \text{H} \quad \text{:}\ddot{\text{Cl}}\text{:} \\    \quad   \\  \text{H}-\text{C}-\text{C}-\ddot{\text{F}}\text{:} \\    \quad   \\  \text{H} \quad \text{:}\ddot{\text{Cl}}\text{:}  \end{array}  $

Chemistry in Context 6<sup>th</sup> Edition, ACS, McGraw-Hill

Table 2.8

Two Important Hydrofluorocarbons

HFC-125	HFC-32
$\text{C}_2\text{HF}_5$ pentafluoroethane	$\text{CH}_2\text{F}_2$ difluoromethane
$  \begin{array}{c}  \text{:}\ddot{\text{F}}\text{:} \quad \text{:}\ddot{\text{F}}\text{:} \\    \quad   \\  \text{H}-\text{C}-\text{C}-\ddot{\text{F}}\text{:} \\    \quad   \\  \text{:}\ddot{\text{F}}\text{:} \quad \text{:}\ddot{\text{F}}\text{:}  \end{array}  $	$  \begin{array}{c}  \text{:}\ddot{\text{F}}\text{:} \\    \\  \text{H}-\text{C}-\ddot{\text{F}}\text{:} \\    \\  \text{H}  \end{array}  $

## Naming of HCFC's and Halons

<b>CFC</b>	<b>ChloroFluoroCarbons</b>
<b>0</b>	Number of double bonds, omitted if zero
<b>1</b>	Number of carbon atoms minus 1, omitted if zero
<b>2</b>	Number of hydrogen atoms plus 1
<b>3</b>	Number of fluorine atoms
<b>4</b>	Number of chlorine atoms replaced by bromine, always used with prefix "b" (b1, b2), omitted if zero
<b>a</b>	Added to identify isomers, the isomer without suffix always has the smallest mass difference on each carbon atom. If there are more isomers the suffix is counting from a - z, omitted if only one isomer exists

# Global warming

**Table 3.3**

**Global Warming Potential for Three Greenhouse Gases**

Substance	Global Warming Potential (GWP)*	Tropospheric Abundance (ppm)	
CO <sub>2</sub>	1	385	<b>385</b>
CH <sub>4</sub>	23	1.8	<b>41</b>
N <sub>2</sub> O	296	0.31	<b>92</b>

Net effect

\*GWP values are given for the estimated direct and indirect effects over a 100-year period and are relative to the assigned value of 1 for CO<sub>2</sub>.

**Table 3.2**

**Greenhouse Gases—Concentration Changes and Lifetimes**


	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
Preindustrial concentration (1750)	278 ppm	0.700 ppm	0.270 ppm
2005 concentration	385 ppm	1.75 ppm	0.314 ppm
Average rate of concentration change, 1990–2005	1.5 ppm/year	0.007 ppm/year	0.0008 ppm/year
Global atmospheric lifetime	50–200 years*	12 years	114 years

Nitrous oxide

\*A single value for the atmospheric lifetime of CO<sub>2</sub> is not possible. Different removal mechanisms take place at different rates, leading to variation in atmospheric lifetime.

*Chemistry in Context 6<sup>th</sup> Edition, ACS, McGraw-Hill*

## Global Atmospheric Concentrations of Carbon Dioxide Over Time



Data source: Compilation of 10 underlying datasets. See [www.epa.gov/climatechange/indicators/ghg/ghg-concentrations.html](http://www.epa.gov/climatechange/indicators/ghg/ghg-concentrations.html) for specific information.


For more information, visit U.S. EPA's "Climate Change Indicators in the United States" at [www.epa.gov/climatechange/indicators](http://www.epa.gov/climatechange/indicators).



## Global Atmospheric Concentrations of Methane Over Time

800,000 BC to 2013 AD


1950 to 2013 AD



Data source: Compilation of five underlying datasets. See [www.epa.gov/climatechange/indicators/ghg/ghg-concentrations.html](http://www.epa.gov/climatechange/indicators/ghg/ghg-concentrations.html) for specific information.


For more information, visit U.S. EPA's "Climate Change Indicators in the United States" at [www.epa.gov/climatechange/indicators](http://www.epa.gov/climatechange/indicators).

## Global Atmospheric Concentrations of Nitrous Oxide Over Time



Data source: Compilation of six underlying datasets. See [www.epa.gov/climatechange/indicators/ghg/ghg-concentrations.html](http://www.epa.gov/climatechange/indicators/ghg/ghg-concentrations.html) for specific information.

For more information, visit U.S. EPA's "Climate Change Indicators in the United States" at [www.epa.gov/climatechange/indicators](http://www.epa.gov/climatechange/indicators).




Global temperatures track CO<sub>2</sub> concentrations very closely

**Figure 3.4**

Atmospheric CO<sub>2</sub> concentration (red) and temperature change from the 1950-1980 mean (blue) over 160,000 years (ice core data).

**Comparison to 1950-1980 mean values**




**Figure 3.14**

Infrared spectrum of carbon dioxide.

The letters (b), (c), and (d) refer to the molecular vibrations shown in Figure 3.13.


**N<sub>2</sub> and O<sub>2</sub> have no effect: no dipole moment change on vibration**



**Figure 3.13**

Molecular vibrations in CO<sub>2</sub>.

Each spring represents a C-to-O double bond. Vibrations a and b are stretching vibrations; c and d are bending vibrations.



**Figure 3.15**

Infrared spectrum of water vapor.

## Intergovernmental Panel on Climate Change – 2007 Synthesis Report

[http://www.ipcc.ch/publications\\_and\\_data/ar4/syr/en/contents.html](http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html)

**Table 3.4**

**Judgmental Estimates of Confidence**

Term	Probability That a Result Is True
Virtually certain	> 99%
Very likely	90–99%
Likely	66–90%
Medium likelihood	33–66%
Unlikely	10–33%
Very unlikely	1–10%

Source: *Summary for Policymakers, A Report of Working Group I of the Intergovernmental Panel on Climate Change*, Shanghai: IPCC, January 1, 2001.


*Chemistry in Context 6<sup>th</sup> Edition, ACS, McGraw-Hill*

Summary of how to interpret language for policy makers!

### **Observed changes in climate:**

*“Warming of the climate system is **unequivocal**, as is now evident from observations of increases in global average air and ocean temperatures, widespread melting of snow and ice and rising global average sea level”*

Changes in temperature, sea level and Northern Hemisphere snow cover




**Figure SPM.1.** Observed changes in (a) global average surface temperature; (b) global average sea level from tide gauge (blue) and satellite (red) data and (c) Northern Hemisphere snow cover for March–April. All differences are relative to corresponding averages for the period 1961–1990. Smoothed curves represent decadal averaged values while circles show yearly values. The shaded areas are the uncertainty intervals estimated from a comprehensive analysis of known uncertainties (a and b) and from the time series (c). {Figure 1.1}

[http://www.ipcc.ch/publications\\_and\\_data/ar4/syr/en/contents.html](http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html)

“Observational evidence from **all continents** and most oceans shows that many natural systems are being affected by regional climate changes, particularly temperature increases”

Global and continental temperature change



**Figure SPM.4.** Comparison of observed continental- and global-scale changes in surface temperature with results simulated by climate models using either natural or both natural and anthropogenic forcings. Decadal averages of observations are shown for the period 1906-2005 (black line) plotted against the centre of the decade and relative to the corresponding average for the period 1901-1950. Lines are dashed where spatial coverage is less than 50%. Blue shaded bands show the 5 to 95% range for 19 simulations from five climate models using only the natural forcings due to solar activity and volcanoes. Red shaded bands show the 5 to 95% range for 58 simulations from 14 climate models using both natural and anthropogenic forcings. (Figure 2.5)

[http://www.ipcc.ch/publications\\_and\\_data/ar4/syr/en/contents.html](http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html)

“There is medium confidence that other effects of regional climate change on natural and human environments are emerging, although many are difficult to discern due to adaptation and non-climatic drivers”

### **Causes of change:**

“Global GHG emissions due to human activities have grown since pre-industrial times, with an **increase of 70% between 1970 and 2004**


Global atmospheric concentrations of CO<sub>2</sub>, methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) have increased markedly as a result of human activities since 1750 and now **far exceed pre-industrial values** determined from ice cores spanning many thousands of years

Most of the observed increase in global average temperatures since the mid-20th century is **very likely** due to the observed increase in anthropogenic GHG concentrations. It is **likely** that there has been significant anthropogenic warming over the past 50 years averaged over each continent (except Antarctica)”

[http://www.ipcc.ch/publications\\_and\\_data/ar4/syr/en/contents.html](http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html)



### Global anthropogenic GHG emissions



**Figure SPM.3.** (a) Global annual emissions of anthropogenic GHGs from 1970 to 2004.<sup>5</sup> (b) Share of different anthropogenic GHGs in total emissions in 2004 in terms of carbon dioxide equivalents (CO<sub>2</sub>-eq). (c) Share of different sectors in total anthropogenic GHG emissions in 2004 in terms of CO<sub>2</sub>-eq. (Forestry includes deforestation.) {Figure 2.1}

[http://www.ipcc.ch/publications\\_and\\_data/ar4/syr/en/contents.html](http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html)

**“Advances since the Third Assessment Report show that discernible human influences extend beyond average temperature to other aspects of climate:**

**Human influences have: {2.4}**


- **very likely** contributed to **sea level rise** during the latter half of the 20th century
- **likely** contributed to **changes in wind patterns**, affecting extra-tropical storm tracks and temperature patterns
- **likely** increased temperatures of **extreme** hot nights, cold nights and cold days
- **more likely than not** increased risk of heat waves, area affected by drought since the 1970s and frequency of heavy precipitation events”

**“Anthropogenic warming over the last three decades has likely had a discernible influence at the global scale on observed changes in many physical and biological systems”**

[http://www.ipcc.ch/publications\\_and\\_data/ar4/syr/en/contents.html](http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html)

# Top Greenhouse Gas Emitters by Country


Top 10 Emitters



<http://bit.ly/11SMpjA>

WORLD RESOURCES INSTITUTE

Per Capita Emissions for Top 10 Emitters



<http://bit.ly/11SMpjA>

WORLD RESOURCES INSTITUTE

**Alternative viewpoints on climate change have been expressed.**

*See for example:*

<http://climatechangereconsidered.org/climate-change-reconsidered-ii-physical-science/>

[https://www.youtube.com/watch?v=dSVkSCN\\_hLQ](https://www.youtube.com/watch?v=dSVkSCN_hLQ)

*...but always bear in mind the source of these ideas and consider carefully whether you think this is unbiased and genuine.*