

## Hydrogen Storage

DOI: 10.1002/anie.200601434

**Storing and Releasing Hydrogen with a Redox Switch\*\****Simon K. Brayshaw, Jennifer C. Green, Nilay Hazari, J. Scott McIndoe, Frank Marken, Paul R. Raithby, and Andrew S. Weller\**

Hydrogen storage is an area of intense current interest, which is driven by the requirements for reversible storage of dihydrogen for future energy applications.<sup>[1]</sup> Chemisorption of dihydrogen in the form of light metal hydrides is one approach to the problem of storing large amounts of H<sub>2</sub> reversibly.<sup>[2]</sup> Physisorption of dihydrogen onto porous network materials such as activated carbon, metal–organic framework (MOF) materials,<sup>[3]</sup> or related systems<sup>[4,5]</sup> presents an alternative method, and MOF systems that incorporate open metal sites (such as Ni<sup>2+</sup>, Cu<sup>2+</sup>, and Mg<sup>2+</sup>)<sup>[5–8]</sup> have recently attracted attention because of the stronger binding of H<sub>2</sub> with the metal–ligand framework.<sup>[8]</sup> Although significant and elegant advances have been made in both areas,

[\*] Dr. S. K. Brayshaw, Dr. F. Marken, Prof. P. R. Raithby, Dr. A. S. Weller  
Department of Chemistry  
University of Bath  
Bath, BA2 7AY (UK)  
Fax: (+44) 1225-386-231  
E-mail: a.s.weller@bath.ac.uk  
Prof. J. C. Green, N. Hazari  
Inorganic Chemistry Laboratory  
South Parks Road, Oxford OX1 3QR (UK)  
Prof. J. S. McIndoe  
Department of Chemistry  
University of Victoria  
PO Box 3065, Victoria, BC V8W3V6 (Canada)

[\*\*] We thank the EPSRC (grant GR/T10169), the Royal Society, and the Rhodes Trust for support, Dr Damien Murphy (Cardiff University) for providing the EPR spectra, and the Oxford Supercomputing Centre for computational facilities.



Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

challenges still remain. Metal hydrides require temperature cycling to promote loss of hydrogen, as hydrogen is chemisorbed (binding energies greater than  $100 \text{ kJ mol}^{-1}$ ). Although porous networks show reversible  $\text{H}_2$  uptake without temperature cycling, they generally require an overpressure to retain  $\text{H}_2$ , because reducing the partial pressure results in loss of  $\text{H}_2$  (i.e., there is no hysteresis between the absorption and desorption isotherms); this is because dihydrogen is physisorbed<sup>[9]</sup> (binding energies less than  $10 \text{ kJ mol}^{-1}$ ). Although these MOF materials can uptake  $\text{H}_2$  very well at  $77 \text{ K}$ ,<sup>[7,10]</sup> adsorption drops dramatically (less than 0.3%) at ambient temperature. MOF systems that show hysteresis have been described recently. These materials retain  $\text{H}_2$  at reduced pressure (ca. 0.01 atm); however, they only operate at  $77 \text{ K}$ .<sup>[11]</sup> In view of the current state of this research, a material that is both a true store of hydrogen under ambient conditions of temperature and pressure and that releases the stored  $\text{H}_2$  without significant external energy input would be of fundamental interest.

We report herein what we believe to be a novel mechanism for hydrogen storage and subsequent release. The system consists of molecular clusters in which  $\text{H}_2$  is adsorbed at binding energies intermediate between the accepted limits for chemi- and physisorption. These clusters retain the stored dihydrogen even at low pressure, which makes them genuine hydrogen stores, but they can release  $\text{H}_2$  rapidly (on the millisecond timescale) when triggered by a simple redox process. Moreover, these processes occur under the attractive conditions of  $298 \text{ K}$  and  $1 \text{ atm H}_2$  and can be triggered either chemically or electrochemically. Although cluster species with hydride ligands are common,<sup>[12–17]</sup> the reversible uptake and release of  $\text{H}_2$  on molecular clusters is rare and examples have generally required relatively forcing conditions (vacuum, heat, photolysis, or addition of an external hydrogen acceptor) to release the stored  $\text{H}_2$ .<sup>[18–20]</sup>

We recently reported the synthesis of a new class of rhodium clusters, exemplified by  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}][\text{BAR}^{\text{F}}_4]_2$  (**A**) ( $[\text{BAR}^{\text{F}}_4] = [\text{B}\{\text{C}_6\text{H}_3(\text{CF}_3)_2\}_4]^-$ ; Cy = cyclohexyl), which have molecular geometries that feature an octahedral hexarhodium core and a trialkyl phosphine ligand radiating from each metal vertex,<sup>[19,21,22]</sup> thus shrouding the metal core with a hydrocarbon shell. These dicationic clusters are completed by twelve hydrogen atoms that bridge each Rh–Rh edge in the octahedron. Calculations<sup>[21]</sup> show that there is only a modest energy gap (0.3 eV) between the highest occupied molecular orbital (HOMO) and a degenerate set of two low-lying unoccupied molecular orbitals (LUMOs, Figure 1). Thus, these clusters are predisposed to receive four extra electrons. We have shown that these electrons can come from two  $\text{H}_2$  molecules (i.e. two bonding electron pairs) and that the addition of  $\text{H}_2$  to **A** affords the 16-hydrogen-atom cluster  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{16}][\text{BAR}^{\text{F}}_4]_2$  (**B**), in which the HOMO–LUMO energy gap is much larger (1.3 eV). The uptake of two molecules of  $\text{H}_2$  is facile (both in the solution and the solid state) and can be reversed by placing **B** under a dynamic vacuum (ca.  $6 \times 10^{-6} \text{ atm}$ ,  $t_{1/2} = 48 \text{ h}$ ). Under ambient conditions (1 atm argon,  $298 \text{ K}$ ), cluster **B** retains the two absorbed  $\text{H}_2$  molecules for weeks with no partial pressure of  $\text{H}_2$ .<sup>[21]</sup>






Figure 1. Frontier molecular orbital energy diagram for **A** and **B**.

Inspection of the MO diagram for **B** suggests that the addition of one electron would result in the population of a high-lying orbital and that this could destabilize the cluster, which would result in  $\text{H}_2$  loss to reestablish a large HOMO–LUMO gap. Exactly this process does in fact occur, and it can be readily followed chemically and electrochemically by using standard spectroscopic and electrochemical techniques. On a preparative scale the addition of the reducing agent  $[\text{Cr}(\eta^6\text{-C}_6\text{H}_6)_2]$  ( $E_{1/2} = -1.15 \text{ V}$  relative to ferrocene/ferrocenium) to a solution of **B** in  $\text{CH}_2\text{Cl}_2$  results in a paramagnetic species that we have unequivocally characterized as  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{14}][\text{BAR}^{\text{F}}_4]$  (**C**) by following the reaction with electrospray ionisation mass spectrometry (ESI-MS). These spectra show the clean formation of **C**, a monocation with fourteen hydrides associated with a  $\{\text{Rh}_6(\text{PCy}_3)_6\}$  core (Figure 2). As might be expected,  $^1\text{H NMR}$  spectroscopy of paramagnetic **C** is relatively uninformative and the hydride signals are not observed. However, H/D exchange experiments using  $\text{D}_2$  show the formation of  $\text{HD}_{(g)}$  and  $\text{H}_2_{(g)}$ , which is consistent with accessible dihydrogen ligands on the cluster surface,<sup>[23]</sup> as are theoretical calculations (see below). Attempts to isolate **C** in pure form on a preparative scale by repeated recrystallization to remove the reductant always led to the loss of  $\text{H}_2$  and isolation of the twelve-hydride monocationic cluster  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}][\text{BAR}^{\text{F}}_4]$  (**D**). The “uncharged-with- $\text{H}_2$ ” starting material **A** is simply returned by oxidation of **C** with  $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2][\text{PF}_6]$  ( $E_{1/2} = 0.00 \text{ V}$ ), as monitored by NMR spectroscopy and ESI-MS. Oxidation of **C** thus results in the elimination of a further equivalent of  $\text{H}_2$ , and a complete hydrogen store and release cycle is completed in solution for two molecules of  $\text{H}_2$  per cluster by a simple redox process.

This redox-triggered release of  $\text{H}_2$  can also be followed electrochemically. Figure 3 shows the cyclic voltammogram of **B** (in  $\text{CH}_2\text{Cl}_2$ ,  $[\text{NBu}_4][\text{BAR}^{\text{F}}_4]$  electrolyte, glassy carbon



**Figure 2.** Redox triggered hydrogen storage/release cycle. i)  $[\text{Cr}(\eta^6\text{-C}_6\text{H}_6)_2]$ ; ii)  $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2][\text{PF}_6]$ ; iii)  $\text{H}_2$ , glassy carbon electrode. Insets show ESI-MS data (positive-ion mode) of cluster species for each step with the observed isotope pattern overlaid with that predicted (gray bars). See the Supporting Information for full spectra.




**Figure 3.** Top: Cyclic voltammogram of  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}]^{2+}$  (**A**) (which is also identical to those of **C** and **D**) showing the three electrochemical processes (two reversible reductions and one quasireversible oxidation). Bottom: Cyclic voltammogram of **B** showing the irreversible reduction (i) and subsequent electrochemistry associated with **A** (ii–iv). See the Supporting Information for full experimental details.

electrode, sweep rate  $100 \text{ mV s}^{-1}$ ). Cluster **B** undergoes an irreversible one-electron reduction at  $E_{1/2} = -1.00 \text{ V}$ , relative to ferrocene/ferrocenium, which is consistent with the irreversible elimination of  $\text{H}_2$ . By progressively increasing the scan rate to  $10 \text{ V s}^{-1}$  and observing the back oxidation, a half-life for the unstable reduced species of  $170 \text{ ms}$  ( $k = 4 \text{ s}^{-1}$  at  $293 \text{ K}$ ) was obtained. There are also three reversible one-electron processes at  $E_{1/2} = -1.51, -0.59, \text{ and } +0.44 \text{ V}$ , which are also observed for **A**<sup>[24]</sup> and **D**, the latter related to **C** by the simple loss of one  $\text{H}_2$  molecule. Presumably under electrochemical conditions, **C** undergoes facile loss of  $\text{H}_2$  to form **D**, and this suggestion is supported by the small calculated

energy difference between these two species ( $0.12 \text{ eV}$ , see below) and our observation that  $\text{H}_2$  is slowly lost from **C** in bulk solution. Cluster **D** can be formed independently in excellent yield by chemical reduction of **A** with  $[\text{Cr}(\eta^6\text{-C}_6\text{H}_6)_2]$  (see the Supporting Information for full characterization, including the X-ray crystal structure). Overall, the electrochemical behavior of **B** can be summarized as the irreversible loss of two molecules of  $\text{H}_2$  upon reduction at the carbon electrode to give **D**, which is directly related to the starting cluster **A** by simple oxidation. Although there is a subtle change in the order of events

between the bulk chemical and electrochemical processes, the overall redox hydrogen storage/release cycle holds for both: the reduction of **B** and subsequent reoxidation to afford **A** is effectively quantitative, repeatable, and involves two equivalents of  $\text{H}_2$  per cluster per cycle.

Full geometry optimizations and calculations of the relative Gibbs free energies were performed using density functional theory (DFT) (see the Supporting Information for details) on the model system  $[\text{Rh}_6(\text{PH}_3)_6\text{H}_x]^{n+}$  ( $x = 12, 14, 16; n = 1, 2$ ). The relative energies of the possible intermediates in this cycle (Gibbs free energy at  $298.15 \text{ K}$ , taking into account solvent effects ( $\text{CH}_2\text{Cl}_2$ ), zero-point energy, and entropy) are shown in Figure 4. The addition of two molecules of  $\text{H}_2$  to  $[\text{H}_{12}]^{2+}$  (**A**) to give  $[\text{H}_{16}]^{2+}$  (**B**) is an overall exothermic process, which is consistent with the experimentally observed formation of **B** on exposure to  $\text{H}_2$  (the “charged-with- $\text{H}_2$ ”



**Figure 4.** Calculated energies (eV) relative to **A** ( $[\text{H}_{12}]^{2+}$ ) for the cluster hydride species in the storage/release cycle. Molecular species in bold face were observed spectroscopically and in some cases [**A**, **B**,<sup>[21]</sup> and **D** (this work)] determined by X-ray diffraction. Energies quoted take into account the Gibbs free energy of gaseous  $\text{H}_2$  where appropriate.  $\text{PCy}_3$  was replaced by  $\text{PH}_3$  in the model systems.

state). A reasonable supposition is that this proceeds via an intermediate dicationic species with fourteen hydrides, namely  $[\text{Rh}_6(\text{PH}_3)_6\text{H}_{14}]^{2+}$  ( $[\text{H}_{14}]^{2+}$ ). The average interaction energy of the two extra  $\text{H}_2$  molecules with the cluster was determined to be 0.62 eV (60 kJ mol<sup>-1</sup>) per dihydrogen. This is comparable with the binding energy both calculated and experimentally determined for dihydrogen bound to a single metal center (60–100 kJ mol<sup>-1</sup>),<sup>[23]</sup> and DFT calculations and H/D exchange experiments both strongly suggest two dihydrogen ligands on the cluster surface of this sixteen-hydride species.<sup>[21]</sup> Addition of one electron to  $[\text{H}_{16}]^{2+}$  results in  $[\text{Rh}_6(\text{P}(\text{PH}_3)_3)_6\text{H}_{16}]^+$  ( $[\text{H}_{16}]^+$ ), which is unstable towards  $\text{H}_2$  loss to afford  $[\text{H}_{14}]^+$  and  $\text{H}_2$ . The calculated structure of  $[\text{H}_{14}]^+$  reveals a relatively close H–H contact on the cluster surface (1.06 Å), which is suggestive of a dihydrogen ligand. Oxidation of  $[\text{H}_{14}]^+$  yields  $[\text{Rh}_6(\text{PH}_3)_6\text{H}_{14}]^{2+}$  ( $[\text{H}_{14}]^{2+}$ ). Although this species is marginally lower in energy than  $[\text{H}_{12}]^{2+}$ , there must be only a small barrier towards  $\text{H}_2$  loss, which can be overcome simply by loss of hydrogen to the bulk solution. The barrier to loss of  $\text{H}_2$  from  $[\text{H}_{14}]^+$  must be larger, as cluster **C** is observed experimentally in bulk solution by ESI-MS. Although the precise mechanism of  $\text{H}_2$  loss is currently uncertain, the correlation between experiment and theory is reasonable, especially given that the approximations involved in using  $\text{PH}_3$  instead of  $\text{PCy}_3$  mean that kinetic factors towards  $\text{H}_2$  loss may not be modeled accurately. Importantly, the experimental observations show unequivocally that the overall hydrogen storage/release cycle (**A** ⇌ **B**) involves two molecules of  $\text{H}_2$  per cluster irrespective of the precise mechanistic details.

The hydrogen storage capacity of **A** (0.1% w/w) is modest, as the cluster and associated anions have high molecular mass, and this value is clearly not sufficient for practical purposes (the US Department of Energy 2010 target is 6% w/w<sup>[25]</sup>). However, the ability to store  $\text{H}_2$  without a partial pressure of  $\text{H}_2$  (and even under dynamic vacuum) at 298 K, coupled with the very rapid redox-promoted release of  $\text{H}_2$ , is unique. These attractive features result from the electronic and steric properties of the metal cluster, namely: 1) two low-lying unoccupied molecular orbitals available for the uptake of bonding electron pairs from  $\text{H}_2$ ; 2) the resulting calculated binding energy of  $\text{H}_2$  to the cluster of 60 kJ mol<sup>-1</sup> (lying at a value intermediate between physis- and chemisorption regimes); and 3) the kinetic stabilization afforded against facile hydrogen loss by the trialkyl phosphine groups. These results suggest that future strategies involving redox-switchable metal clusters in hydrogen storage materials or hydrogen spillover systems<sup>[26]</sup> could well prove worthwhile.

Received: April 11, 2006  
Published online: August 4, 2006

**Keywords:** cluster compounds · density functional calculations · electrochemistry · hydrogen storage

- [3] J. L. C. Rowsell, O. M. Yaghi, *Angew. Chem.* **2005**, *117*, 4748; *Angew. Chem. Int. Ed.* **2005**, *44*, 4670.
- [4] N. B. McKeown, B. Gahnem, K. J. Msayib, P. M. Budd, C. E. Tattershall, K. Mahmood, S. Tan, D. Book, H. W. Langmi, A. Walton, *Angew. Chem.* **2006**, *118*, 1836; *Angew. Chem. Int. Ed.* **2006**, *45*, 1804.
- [5] S. S. Kaye, J. R. Long, *J. Am. Chem. Soc.* **2005**, *127*, 6506.
- [6] P. M. Forster, J. Eckert, J. S. Chang, S. E. Park, F. G. , A. K. Cheetham, *J. Am. Chem. Soc.* **2003**, *125*, 1309.
- [7] B. Chen, N. W. Ockwig, A. R. Millward, D. S. Contreras, O. M. Yaghi, *Angew. Chem.* **2005**, *117*, 4823; *Angew. Chem. Int. Ed.* **2005**, *44*, 4745.
- [8] M. Dinca, J. R. Long, *J. Am. Chem. Soc.* **2005**, *127*, 9376.
- [9] J. L. C. Roswell, E. C. Spencer, J. Eckert, J. A. K. Howard, O. M. Yaghi, *Science* **2005**, *309*, 1350.
- [10] a) G. Férey, M. Lacroche, C. Serre, F. Millange, T. Loiseau, A. Percheron-Guegan, *Chem. Commun.* **2003**, 2976; b) G. Férey, C. Mellot-Draznieks, C. Serre, F. Millange, J. Dutour, S. Surlblé, I. Margiolaki, *Science* **2005**, *309*, 2040; c) A. G. Wong-Foy, A. J. Matzger, O. M. Yaghi, *J. Am. Chem. Soc.* **2006**, *128*, 3494.
- [11] X. Zhao, B. Xiao, A. J. Fletcher, K. M. Thomas, D. Bradshaw, M. J. Rosseinsky, *Science* **2004**, *306*, 1012.
- [12] J. M. Bemis, L. F. Dahl, *J. Am. Chem. Soc.* **1997**, *119*, 4545.
- [13] M. H. Chisholm, K. S. Kramer, *Chem. Commun.* **1996**, 1331.
- [14] L. F. Chen, F. A. Cotton, W. T. Klooster, T. F. Koetzle, *J. Am. Chem. Soc.* **1997**, *119*, 12175.
- [15] P. Braunstein, L. A. Oro, P. R. Raithby, *Metal Clusters in Chemistry*, Wiley-VCH, Weinheim, **1999**.
- [16] H. Suzuki, T. Kakigano, K. Tada, M. Igarashi, K. Matsubara, A. Inagaki, M. Oshima, T. Takao, *Bull. Chem. Soc. Jpn.* **2005**, *78*, 67.
- [17] T. H. Lemmen, K. Folting, J. C. Huffman, K. G. Caulton, *J. Am. Chem. Soc.* **1985**, *107*, 7774.
- [18] R. D. Adams, B. Captain, M. D. Smith, *Angew. Chem.* **2006**, *118*, 1127; *Angew. Chem. Int. Ed.* **2006**, *45*, 1109.
- [19] S. K. Brayshaw, M. J. Ingleson, J. C. Green, P. R. Raithby, G. Kociok-Köhn, J. S. McIndoe, A. S. Weller, *Angew. Chem.* **2005**, *117*, 7035; *Angew. Chem. Int. Ed.* **2005**, *44*, 6875.
- [20] R. D. Adams, B. Captain, *Angew. Chem.* **2005**, *117*, 2587; *Angew. Chem. Int. Ed.* **2005**, *44*, 2531.
- [21] S. K. Brayshaw, M. J. Ingleson, G. Kociok-Köhn, J. C. Green, J. S. McIndoe, P. R. Raithby, A. S. Weller, *J. Am. Chem. Soc.* **2006**, *128*, 6247.
- [22] M. J. Ingleson, M. F. Mahon, P. R. Raithby, A. S. Weller, *J. Am. Chem. Soc.* **2004**, *126*, 4784.
- [23] G. J. Kubas, *Metal Dihydrogen and σ-Bond Complexes*, Kluwer Academic/Plenum Publishers, New York, **2001**.
- [24] These redox events correspond to  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}]^0/[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}]^+$   $E_{1/2} = -1.51$ ,  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}]^{2+}/[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}]^{2+}$   $E_{1/2} = -0.59$ , and the quasireversible  $[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}]^{2+}/[\text{Rh}_6(\text{PCy}_3)_6\text{H}_{12}]^{3+}$   $E_{1/2} = +0.44$  V. Full characterization (including X-ray crystal structures, EPR studies, magnetic properties, and DFT calculations) of these species will appear in a full paper. For a recent example of related electrochemical behavior in early-transition-metal clusters, see: E. J. Welch, C. L. Yu, N. R. M. Crawford, J. R. Long, *Angew. Chem.* **2005**, *117*, 2605; *Angew. Chem. Int. Ed.* **2005**, *44*, 2549.
- [25] Basic Research Needs for the Hydrogen Economy; [http://www.eere.energy.gov/hydrogenandfuelcells/pdfs/hydrogen\\_posture\\_plan.pdf](http://www.eere.energy.gov/hydrogenandfuelcells/pdfs/hydrogen_posture_plan.pdf).
- [26] Y. Li, R. T. Yang, *J. Am. Chem. Soc.* **2006**, *128*, 726.

[1] L. Schlapbach, A. Züttel, *Nature* **2001**, *414*, 353.

[2] F. Schüth, B. Bogdanovic, M. Felderhoff, *Chem. Commun.* **2004**, 2249.