frontiers in NEUROANATOMY

Genotype-specific effects of MECP2 loss-of-function on morphology of Layer V pyramidal neurons in heterozygous female Rett mice

Leslie Rietveld, David P Stuss, David McPhee and Kerry R. Delaney

Journal Name:	Frontiers in Neuroanatomy
ISSN:	1662-5129
Article type:	Original Research Article
First received on:	13 Dec 2014
Frontiers website link:	www.frontiersin.org

Genotype-specific effects of MECP2 loss-of-function on morphology of Layer V pyramidal neurons in heterozygous female Rett mice

Leslie Rietveld, David P. Stuss¹, David McPhee and Kerry R. Delaney*

Dept. of Biology, University of Victoria, Victoria BC Canada V8W2Y2; ¹Present address: Dept of Chemistry and Geoscience, Camosun College, Victoria BC Canada V8P5J2

Corresponding author

K. R. Delaney, Dept. of Biology PO Box 1700 STN CSC Victoria BC Canada V8W2Y2 kdelaney@uvic.ca 1 250 721 7091

ABSTRACT

Rett Syndrome (RTT) is a progressive neurological disorder primarily caused by mutations in the Xlinked gene methyl-CpG-binding protein 2 (MECP2). The heterozygous female RTT mouse brain consists of mosaic of neurons expressing both wildtype MECP2 (Mecp2+) and mutant MECP2 (Mecp2-). 3-dimensional morphological analysis was performed on individually genotyped layer V pyramidal neurons in the primary motor cortex of heterozygous $(Mecp2^{+/-})$ and wild-type $(Mecp2^{+/+})$ female mice (>6 mo.) from the $Mecp2^{tm1.1Jae}$ line. Comparing basal dendrite morphology, soma and nuclear size of Mecp2+ to Mecp2- neurons reveals a significant cell autonomous, genotype specific effect of Mecp2. Mecp2- neurons have 15% less total basal dendritic length, predominantly in the region 70-130 µm from the cell body and on average 3 fewer branch points, specifically loss in the 2nd and 3rd branch orders. Soma and nuclear areas of neurons of mice were analyzed across a range of ages (5-21 mo.) and Xchromosome inactivation (XCI) ratios (12-56%). On average, Mecp2- somata and nuclei were 15% and 13% smaller than Mecp2+ neurons respectively. In most respects branching morphology of neurons in wild-type brains (Mecp2+/+) was not distinguishable from Mecp2+ but somata and nuclei of Mecp2+/+ neurons were larger than those of Mecp2+ neurons. These data reveal cell autonomous effects of MECP2 mutation on dendritic morphology, but also suggest non-cell autonomous effects with respect to cell size. Mecp2+ and Mecp2- neuron sizes were not correlated with age, but were correlated with XCI ratio. Unexpectedly the Mecp2- neurons were smallest in brains where the XCI ratio was highly skewed towards Mecp2+, i.e. wild-type. This raises the possibility of cell non-autonomous effects that act through mechanisms other than globally secreted factors; perhaps competition for synaptic connections influences cell size and morphology in Rett mosaic brains.

INTRODUCTION

Rett Syndrome (RTT) is a postnatal neurological disorder that was first characterized by Andreas Rett in 1966 (Rett, 1966). It affects approximately one in 10,000 live female births (Laurvick *et al.*, 2006). Typical RTT patients are diagnosed after a period of normal development (6-18 months of age) followed by a period of regression marked by four main criteria: partial or complete loss of purposeful hand movements, development of gait abnormalities, loss of verbal language skills and the presence of stereotypic hand movements (Neul *et al.*, 2010). The majority of females with RTT are heterozygous for a mutation in the X-linked gene Methyl-CpG-binding Protein 2 (*MECP2*) (Amir *et al.*, 1999). Normally, MECP2 expression levels are low prenatally, but increase after birth during the final stages of neurogenesis (Balmer *et al.*, 2003; Kishi and Macklis, 2004; Skene *et al.*, 2010), supporting the observation that MECP2 contributes to neuronal dendritic maturation and synaptogenesis (Armstrong *et al.*, 1995; Kishi and Macklis, 2004; Fukuda *et al.*, 2005). Although over 1000 mutations have been characterized along the entire length of the gene, including nonsense, missense, frameshift, and large truncation mutations (Amir *et al.*, 2000; Weaving *et al.*, 2005; Philippe *et al.*, 2006; Cuddapah *et al.*, 2014), 65% of RTT cases are caused by 8 common missense mutations at CpG dinucleotides (Miltenberger-Miltenyl and Laccconi, 2003).

The protein encoded by the *MECP2* gene has five key domains including a highly conserved methyl-CpG binding domain (MDB) characteristic of its protein family, a transcriptional repressor domain, which interacts with histone deacetylases 1 and 2 to promote chromatin condensation, and a C-terminal domain, which contributes to DNA-binding (Chandler, *et al.*, 1999). Phenotype/genotype studies have found that early N-terminal truncation mutations that affect the MBD are correlated with a more severe phenotype than late C-terminal truncation mutations (Zappella *et al.*, 2001; Charman *et al.*, 2005, Cuddapah *et al.*, 2014). The *Mecp2^{tm1.1Jae}* mice used in this study have exon 3 deleted, which comprises most of the MBD. The resulting translated mutant protein may be partly functional (Stuss *et al.*, 2013), leading to a milder phenotype in these animals than has been observed in total knockouts (Belichenko *et al.*, 2008), but still resulting in severe neuropathic symptoms similar to the human condition (Chen *et al.*, 2001).

MECP2 is found in many tissue types, but it is most abundant in the brain, with expression levels in neuronal nuclei ten times higher than in glia (Skene *et al.*, 2010). *MECP2* is located on the Xchromosome and therefore it is affected by X-chromosome inactivation (XCI) (Adler *et al.*, 1995; Ross *et al.*, 2005). XCI occurs in early embryogenesis (gastrulation in humans), randomly inactivating either the maternal or paternal X-chromosome in each cell and passing this status on to all future progeny (Bermejo-Alvarez *et al.*, 2012). XCI therefore results in two populations of neurons in RTT females: those expressing wild-type MECP2 (Mecp2+), and those that lack fully functional MECP2 (Mecp2-) Although the majority of human patients with typical RTT have nearly balanced XCI ratios (Shahbazian *et al.*, 2002), the rate of skewed XCI ratios is nonetheless higher in patients with X-linked disorders such as RTT than in the general population. When skewing occurs it usually favours wild-type over mutant cells (Belmont, 1996; Puck and Willard, 1998; Brown and Robinson, 2000) and XCI ratios follow a similar pattern in mice (Young and Zoghbi, 2004).

RTT patients deficient in MECP2 have reduced grey matter volume in the frontal and temporal lobes, caudate nucleus, thalamus, midbrain, and cerebellum (Reiss *et al.*, 1993; Subramaniam *et al.*, 1997). Volume reductions in the frontal and temporal lobes have been found to be predictive of phenotype severity in RTT (Carter *et al.*, 2008). Neurodegeneration has not been observed in forebrain (Armstrong 1995), and the symptoms of RTT in transgenic mice can be reversed even in adulthood (Luikenhuis *et al.*, 2004; Guy *et al.*, 2007) suggesting that RTT is caused by a defect in neurological function rather than by neuronal damage (McGraw *et al.*, 2011). Morphological studies performed on post-mortem tissue from both RTT patients and mouse models have revealed that the reduction in brain volume is accompanied by increased neuron density associated with reductions in neuronal cell body size (Bauman *et al.*, 1995; Chen *et al.*, 2001). It has also been found that RTT patients have reduced dendritic arborization and spine density of Layer II/III and V/VI pyramidal cells in the frontal and temporal areas, including the motor cortex (Belichenko, 1994; 1997). Similar reductions in spine density have been reported for hippocampal CA1 neurons of RTT patients (Chapleau *et al.*, 2009).

Most animal studies have used Mecp2 mutant male mice $(Mecp2^{-5y})$, since their juvenile symptom onset and rapid progression mirrors RTT patients and shortens the time required for experimental studies. As well, since $Mecp2^{-5y}$ mice express mutant MECP2 in all cells, variations in XCI ratio between individuals are not a potential complication. However, these studies may not address distinct effects that occur in the mosaic brain environment of heterozygous females. $Mecp2^{+5/2}$ mice display the same RTTlike neuropathological and motor abnormalities as $Mecp2^{-5/2}$ mice, but these symptoms are normally only apparent after 4-6 months of age, compared to 3-5 weeks for male mice, with some female mice even living a normal lifespan (Guy *et al.*, 2001; 2007; Chen *et al.*, 2001; Bissonnette *et al.*, 2006; 2007; Abdala *et al.*, 2010).However, progressive motor deficits, reduced anxiety, apnea, and weight gain have been described in $Mecp2^{+5/2}$ mice as young as 3 to 4 weeks old (Santos *et al.*, 2007; Samaco *et al.*, 2009; 2012), which are largely pre-symptomatic for the more readily observed motor abnormalities.

The mosaic brain environment of the female $Mecp2^{+/-}$ mouse prompts the following question: are the cellular phenotypes of Mecp2+ and Mecp2- neurons determined by their individual genotype (cell autonomy) or by the surrounding environment, including neural and non-neuronal cells (non-cell-autonomy), or both?

Our laboratory recently investigated the morphology of pyramidal neurons in the primary motor cortex of Mecp2^{tm1.1Jae} male mice using the YFP-H transgenic mouse line (Feng et al., 2000) to visualize morphology. We found that $Mecp2^{-/y}$ basal and apical dendritic arbours are reduced in length, and that Mecp2^{-/y} spine density is selectively reduced on the apical tuft and on oblique apical dendrites (Stuss et al., 2012). The aim of the current study is to investigate how a mutation in *Mecp2* affects the morphology of both neuronal genotypes in a female $Mecp2^{+/2}$ brain. We examined the morphology of basal dendrites, as well as the sizes of soma and nuclei in Layer V pyramidal neurons of the primary motor cortex, using single-cell microinjection of Alexa Fluor 594 with DAPI and Neurotrace[™] counterstains. Neuronal genotypes were determined by immunohistochemistry. For this study Mecp2- is used to indicate neurons within the brain of Mecp2^{+/-} animals that express mutant MECP2, Mecp2+ refers to neurons expressing wild-type MECP2 and Mecp2+/+ denotes neurons in wild-type female brain. Reduced branching was observed primarily within 30-70 um of the soma in Mecp2- neurons. In most respects dendritic structures of Mecp2+ neurons were not distinguishable from Mecp2+/+ neurons, indicating predominantly cellautonomous effects of the mutation on dendrite branching. Within-animal comparisons showed that soma and nuclear sizes of Mecp2- neurons were reduced compared to Mecp2+ neurons, consistent with cellautonomous effects. Between-animal comparisons revealed that Mecp2+/+ somata were larger than Mecp2+, suggesting an additional non cell-autonomous effect. Intriguingly, the ratio of Mecp2- to Mecp2+ soma and nuclear size was negatively correlated with XCI. The difference in size was greatest when the proportion of Mecp2- neurons was low (highly skewed XCI), which may indicate an additional negative environmental effect for Mecp2- neurons when they are in the minority or a reduced ability of Mecp2- neurons to benefit from a predominantly wild-type brain environment.

METHODS

Experimental Animals

Heterozygous $Mecp2^{+/-}$ female mice ($Mecp2^{tm1.1Jae}$ /Mmcd) (MMRRC, UC Davis) (Chen *et al.*, 2001) were maintained on a C57BL/6 background and their wildtype littermates ($Mecp2^{+/+}$) were used as controls for soma and nuclear size analysis. $Mecp2^{+/-}$ females were crossed with homozygous male YFP-H mice (B6.Cg-Tg(Thy1-YFPH)2Jrs/J) (Feng *et al.*, 2000), which were also maintained on a C57BL/6 background, to generate female offspring heterozygous for both Mecp2 and YFP-H. These offspring ($Mecp2^{+/-}/YFP^{+/-}$) and their wildtype littermates ($Mecp2^{+/+}/YFP^{+/-}$) were used for single-cell microinjection experiments. YFP expression observed with blue excitation confirmed the location of Layer 5 neurons for injection with Alexa 594. All animals were housed in the University of Victoria Animal Care Unit in compliance with the guidelines established by the Canadian Council on Animal Care and Use with approval of protocols by the University Animal Care Committee. Animals were kept at 21° ± 2°C under a 12 hour light/dark cycle with limited environmental enrichment, and were fed standard laboratory diet and water *ad libitum*. *Mecp2* status was determined by genotyping (Stuss *et al.*, 2012).

Prior to tissue extraction the phenotype of the mice was scored as described in Table 1. Mice were then anesthetized and transcardially perfused with 10 ml of room temperature 0.1 M phosphate buffered saline (PBS) pH 7.4 for exsanguination (10 ml/min).

Table 1: Phenotype severity scale for the Jaenisch Rett syndrome mouse model (deletion of exon 3 of *Mecp2*)

Parameter	Severity			
Tremors: observed while standing on the palm	No tremor (0)	Intermittent mild tremor (1)		Continuous tremor or intermittent violent tremor (2)
Hind-clasping: observed when suspended by holding the base of the tail	Legs splay outwards (0)	Legs are drawn inwards slightly (1)	Legs are drawn inwards but do not touch (2)	Legs are drawn inward and touch each other and the body tightly (3)
Dishevelled fur	Clean and shiny coat (0)	Coat dull/ungroomed (1)		Piloerection, dull coat/ungroomed (2)
Activity level: observed when placed on the bench	WT activity level (0)	Slower pace than WT but still active (1)	Still for periods, but can move slowly (2)	No spontaneous movements (3)
Breathing problems	Normal breathing (0)	Short periods of rapid breathing or apnea (1)		Irregular breathing, gasping or panting (2)
Hunched	WT rounded posture (0)	Slightly hunched (1)		Hunched posture (2)

Adapted from (Guy et al., 2007).

For mice prepared for single-cell microinjections, PBS perfusion was followed by 30 ml of 4% paraformaldehyde (4% PFA) in 0.1 M phosphate buffer, pH 6.5, at a rate of 5 ml/min. After removal from the cranium, the brain was post-fixed in 4% PFA, pH 6.5, for 10 minutes at RT. Following this, the brain was removed and placed in a weigh boat for up to 15 minutes before slicing.

For mice prepared for soma and nuclear size analysis (6 to 21 months), the PBS was followed by 10 ml of 4% PFA pH 6.5, (5ml/min) and 20 ml of 0.1M borate buffered 4% PFA, pH 11 at a rate of 2ml/min. After craniotomy, the brain was submerged in 4% PFA, pH 11, for 2 hours at RT. The brain was then immersed in 30% sucrose in PBS and 0.01% sodium azide at 4°C for approximately 48 hours.

Brains were embedded in agar and sliced coronally on a Vibratome Series 1000. The primary motor cortex was identified by the first appearance of the corpus callosum anteriorly and by the joining of the anterior commissure posteriorly (~2 mm). Brains prepared for single cell microinjections were sliced into approximately ten 200 μ m-thick sections and then placed in 24-well plates filled with PBS and 0.01% sodium azide for storage at 4°C. Brains prepared for soma and nuclear size analysis were sliced

into 50 μ m-thick sections and placed in 24-well plates filled with 30% sucrose in PBS and 0.01% sodium azide for storage at 4°C prior to mounting.

For microinjection, the slices were placed in a PBS bath. The cells were visualized using a custom-made epifluorescence microscope equipped with an Olympus UPlanFl 4X/0.13 NA lens and an Olympus LUMPlanFl 40X/0.8 NA water immersion lens. Borosilicate glass pipettes (O.D. 1.5 mm ID 0.86 mm; Sutter Instruments) were prepared with a resistance of 5-10 MΩ when filled with 120 mM KCl (approximately 0.5 µm tip opening) using a micropipette puller (Sutter Instruments P-87). Pipettes were backfilled by dipping in a solution of 3 mM Alexa Fluor 594 hydrazide dye (Invitrogen). Dye fluorescence was excited using a bandpass filter (HQ550/100, Chroma Optical) and visualized through a 624/40 filter (Semrock BrightlineTM) using an analog CCD camera connected to a TV monitor. Cells in Layer V were targeted for filling by locating the YFP+ cells, which were faintly visible using the above filter combination (Figure 1). Individual neurons were targeted using transillumination and the contrast enhancement of the monitor. Cells were impaled and iontophoretically filled with dye using 3 ms square wave, 3 µA current pulses repeated at 0.5 Hz for 5-10 minutes delivered from a digital stimulator (Neuro Data PG4000) through an isolated current source (NeuroData SIU90). Neurons were filled in a sparse distribution along the slice (5-10 cells/hemisphere) to ensure that dendrites did not overlap (Figure 1). Instances in which tissue adhered to the pipette upon tip withdrawal were judged to be a result of poor fixation, typically resulting in dye diffusion out of the cell and/or removal of the nucleus. In these cases the the cell not used for further analysis. Once several cells were filled, the slice was post-fixed in 4% PFA pH 11 for 10 minutes at room temperature and stored in PBS with 0.01% sodium azide at 4°C prior to measurement.

Immunohistochemistry and counterstaining

All immunohistochemistry incubation steps were performed on floating slices at room temperature, on a shaker in the dark, with three PBS washes between each step and 0.01% sodium azide added for overnight incubations. All slices were permeabilized with 1% Triton X-100 in PBS for 2 hours.

Brain slices containing dye filled neurons were immunohistochemically labeled to detect the genotype of individual neurons (Figure 2). Slices were blocked with 5% normal goat serum in PBS overnight, then incubated overnight with monoclonal mouse anti-N terminal Mecp2 (Sigma 4B6) and monoclonal chicken anti-C terminal Mecp2 (Millipore ABE171) primary antibodies prepared at 0.2 µg/ml in 1% normal goat serum and PBS. Primary antibodies were detected with goat anti-mouse conjugated to Alexa Fluor 647 (Invitrogen) and goat anti-chicken conjugated to Alexa Fluor 488 (Invitrogen) secondary antibodies (4 µg/ml) prepared in 1% normal goat serum in PBS and incubated overnight. Slices were counterstained for 10 minutes with 30 nM 4',6-diamidino-2-phenylindole (DAPI) in PBS.

Brain slices prepared for soma and nuclear size analysis (Figure 2) were incubated overnight with monoclonal chicken anti-C terminal Mecp2 (Millipore ABE171) primary antibody prepared at 0.2 μ g/ml in PBS. The primary antibody was detected using goat anti-chicken secondary antibodies conjugated to Alexa Fluor 555 (Invitrogen) prepared at 4 μ g/ml in PBS and incubated overnight. Slices were counterstained with DAPI (30 nM) and the deep-red fluorescent Nissl stain Neurotrace 640/660 (Invitrogen) overnight.

Lipofuscin reduction

Broad-spectrum lipofuscin autofluorescence in older mice interferes with the visibility and quantification of neuronal dendrites (Figure 2). After completing immunohistochemistry and counterstaining protocols, lipofuscin autofluorescence was quenched by treating cortical slices with 5mM CuSO₄ in 50mM ammonium acetate for 10 minutes (Schnell *et al.*, 1999).

Fluorescence microscopy

Slices were mounted on poly-l-lysine coated slides and covered with #1.5 coverslips using Immumount medium (Thermo Scientific). All images for quantification of dendrite structure were obtained using an Olympus UPlanFLN 40X/1.3 NA oil objective. Confocal stacks of single cells microinjected with Alexa Fluor 594 hydrazide were acquired using a 543 nm laser, 610/100 emission filter, 640 X 640 pixel resolution, 0.5 µm steps, 40 µs/pixel dwell, and high laser power. The high voltage, gain, and offset were modified during imaging to optimize visibility of the dendrites and ensure that the end of each dendrite was clearly ascertained. Any dendrites that projected beyond the field of view were captured in an additional adjacent stack. In order to genotype individual neurons, a separate confocal stack centered on the cell body was acquired at 6X zoom with additional channels for detection of the C-terminal (488 nm laser, 515/20 emission) and N-terminal (635 nm laser, 705/100 emission) MECP2 antibodies. Brain slices prepared for soma and nuclear size analysis were viewed on an Olympus IX70 inverted epifluorescence microscope. 5-20 images per animal were captured with a Retiga 2000R digital CCD camera (QImaging, Surrey, BC) with an additional 1.6X post objective magnification. *XCI ratios and soma/nuclear size measurements*

Widefield fluorescence images were obtained from slices of tissue from $30 Mecp2^{+/-}$ animals in one experiment and $11 Mecp2^{+/-}$ and $10 Mecp2^{+/+}$ animals in a second experiment (30-80 cells/ animal). Images were obtained using a UPlanFLN 40X 1.30 NA oil immersion objective (Olympus) focused at 2-3 separate planes, between 10-25 µm below the surface, each separated by approximately 5 µm. Inclusion criteria for measured cells at each focus position were: 1) the DAPI-stained nucleus must be in sharp focus in the focal plane 2) the Neurotrace-stained soma must not be occluded by overlying cells , and 3) the genotype must be clearly identifiable using C-terminal MECP2 antibody staining. Mecp2+ neurons were counted if punctate staining of the C-terminal antibody (colocalized with the DAPI staining) was

visible. Mecp2- neurons were counted if DAPI and diffuse C-terminal antibody staining were visible. Outlines of somata (Neurotrace) and nuclei (DAPI) were manually traced at 8X image magnification using ImageJ (Abràmoff *et al.*, 2004).

Neuronal reconstruction and morphological analysis

82 Mecp2+/+ neurons, 98 MECP2+ and 59 MECP2- neurons were filled and reconstructed (5 animals per genotype, 16-31 cells per animal). In order to minimize variation associated with poor fills or chopped dendrites neurons were excluded if: 1) the genotype could not be determined 2) less than 3 primary dendrites were intact 3) more than half of the total primary dendrites on the cell body were chopped off by the slicing 4) tips of the dendrites were not visible 5) an apical dendrite of at least 75 micrometers length was not visible. If dendrites extended beyond the edge of one field of view additional image stacks were obtained and tiled in 3D using the Volume Integration and Alignment System (VIAS) (Rodriguez *et al.*, 2003). ImageJ was used to adjust the image brightness/contrast and images were saved as 8-bit tiffs. 3D neuronal reconstructions were generated in Neuronstudio (Wearne *et al.*, 2005) using a Cintiq 21q tablet. Neuron tracings were exported in SWC file format, for compatibility with standard morphometric software tools. Multiple morphological parameters were extracted using the freeware applications Neuronstudio, Simple Neurite Tracer (Longair *et al.*, 2011), and L-Measure (Scorcioni *et al.*, 2008). The morphological parameters chosen for analysis in this study are listed in Table 2 and further described in Figure 3 adapted from procedures and conventions described by Scorcioni *et al.*, (2008) and Costa *et al.*, (2010).

Statistical analysis

Statistical analyses were conducted using Prism software (GraphPadTM). Data are presented as means \pm standard error of the mean (SEM). All parameters assessed across Sholl radii or branch orders were analyzed using two-way repeated measures ANOVA with multiple comparisons made using the Bonferroni post-test. Primary dendrite number was assessed using a one-way ANOVA and multiple comparisons were made using the Bonferroni post-test. Direct comparisons of soma and nuclear areas between animals of different genotypes were performed using the nonparametric two-tailed Mann-Whitney U test. Comparisons between cells of different genotypes within the same animal were performed using the paired Wilcoxon signed rank test. Linear regressions were used to compare the slopes and y-intercepts of soma and nuclear area data obtained across ages and XCI ratios. Non-linear regressions were fit to the frequency distributions of soma and nuclear areas and the best-fit values were compared. P-values of ≤ 0.05 were considered significant for all statistical tests.

Table 2 – Morphological parameters analyzed from 3D neuronal reconstructions

Program	Parameter/Description
Neuronstudio	Sholl length (µm) – Total length of all branches per 10 µm Sholl radius
	Cumulative Sholl length (μ m) – Cumulative length of all branches per 10
	μm Sholl radius
	Sholl branch points – Total number of branch points per 10 µm Sholl radius
	Branch length/branch order – Average branch length per branch order
L-Measure	Total number of primary dendrites stemming from the soma
	Total length of all branches per branch order
	Number of branch points per branch order
	Number of branches per branch order
	Number of tips per branch order
	Maximum radial distance (μm) - Euclidean distance between the soma and
	the farthest compartment reached by a dendrite
	Maximum branch order per 10 µm Sholl radius
	Contraction per branch order – ratio between the Euclidean distance and
	branch path distance calculated per branch order
	Partition asymmetry - where T_1 and T_2 are the number of tips of each
	daughter tree. Computed at every branch point and plotted as function of
	branch order

RESULTS

Sholl analysis of basal dendritic arbor of motor cortical Layer V pyramidal neurons of Mecp2+/+, Mecp2+, and Mecp2- neuronal genotypes

Sholl analysis was used to determine how a mutation in *Mecp2* affects the dendritic morphology of Mecp2+ and Mecp2- neurons compared to *Mecp2+/+* neurons. The total dendritic length per 10 μ m Sholl radius was reduced in Mecp2- neurons at distances of 70 to 130 μ m from the soma compared to both Mecp2+/+ and Mecp2+ neurons (2-way repeated measures ANOVA and Bonferroni post-tests, $F_{(62,372)} = 2.33$, p<0.0001; Figure 4A). Mecp2+/+ and Mecp2+ neurons are not different at any Sholl radius (p>0.05). The slopes of the Mecp2+ and Mecp2+/+ curves diverge from that of the Mecp2- curve between 30 and 60 μ m from the cell body, and run parallel at greater distances, indicating that differences in length between the genotypes are initiated in this region and maintained into the distal regions.

To quantify the total differences in dendritic length observed in Figure 4A, cumulative dendritic length was calculated at each successive 10 μ m Sholl radius (Figure 4B). Mecp2- neurons have a significant reduction (15% or 275 μ m) in cumulative dendritic length compared to both other genotypes, while Mecp2+/+ and Mecp2+ are not significantly different (Mecp2- = 1600.6 ± 66.2 μ m, Mecp2+/+ = 1898.5 ± 77.6 μ m, Mecp2+ = 1873.9 ± 66.5 μ m; 2-way repeated measures ANOVA and Bonferroni posttests, F_(44,264) = 4.68, p<0.0001). The maximum radial distance was not different between the genotypes (Mecp2- = 149.7 ± 4.3 μ m, Mecp2+/+ = 159.7 ± 5.9 μ m, Mecp2+ = 155.5 ± 6.2 μ m; one-way ANOVA, F_(2,12) = 0.83, p=0.5) suggesting that although Mecp2- neurons have less length than Mecp2+ neurons, the reach of their dendritic arbour is not significantly lessened.

Branch path distances at successive 10 μ m Sholl radii were compared across genotypes (Figure 5A) to determine whether the cumulative length deficiency of Mecp2- neurons is due to a decreased tortuosity of the branches. No differences were found between the genotypes at any Sholl radii (2-way repeated measures ANOVA and Bonferroni post-tests, $F_{(30,180)} = 1.05$, p=0.4), indicating that the reduction in MECP2- dendritic length is not due to straighter branches.

The number of primary dendrites was compared (Figure 5B) across neuronal genotypes to determine if the differences in length of Mecp2- neurons (Figure 4B) could be accounted for by their having a different number of primary branches. There was no difference in the total number of primary dendrites across all genotypes (one-way ANOVA, $F_{(2,12)} = 0.15$, p=0.9).

Examining the cumulative number of branch points at each successive 10 μ m Sholl radius (Figure 5C) revealed that Mecp2- neurons have an average of three (17%) fewer total branch points than both other genotypes (Mecp2- = 13.7 \pm 0.7, Mecp2+/+ = 16.5 \pm 0.4, Mecp2+ = 16.2 \pm 0.8; t₍₂₄₎=3.51, p< 0.01,

2-way repeated measures ANOVA and Bonferroni post-tests, $F_{(24,144)} = 8.58$, p<0.0001). The slopes of the lines differ between 30 and 60 µm from the soma, suggesting that Mecp2- neurons have fewer branch points in this region.

The reduced number of branch points in Mecp2- cells could be the result of either fewer simple primary trees (low maximum branch order), fewer complex ones (high maximum branch order), or both. In order to determine which was the case, the maximum branch order reached per cell at each successive 10 μ m Sholl radii was compared across the genotypes (Figure 5D). At 60 μ m (Mecp2- = 4 ± 0.20, *Mecp2*+/+ = 5 ± 0.3, Mecp2+ = 5 ± 0.2; t₍₂₄₎=3.43, p<0.05) and 110 μ m (Mecp2- = 4 ± 0.2, *Mecp2*+/+ = 5 ± 0.1, Mecp2+ = 5 ± 0.3; t₍₂₄₎=3.43, p<0.05) from the soma Mecp2- neurons have a lower maximum branch order compared to both other genotypes (2-way repeated measures ANOVA and Bonferroni posttests, F_(2,12) = 1.38, p<0.0001). Since Mecp2+ cells have a relatively high maximum branch order (five) at these distances, this suggests that Mecp2- cells have fewer overall branch points because they have fewer complex branches.

Branch order analysis of Mecp2+/+, Mecp2+, and Mecp2- neuronal genotypes

Branch order analysis was used to determine how a reduction in total branch point number affects the detailed branch structure of Mecp2- neurons (Figure 6). The number of branch points within each branch order was first compared across the neuronal genotypes (Figure 6B). Mecp2- neurons have fewer branch points at the second (Mecp2- = 4.6 ± 0.3 , Mecp2+/+ = 5.3 ± 0.1 , Mecp2+ = 5.1 ± 0.4 ; $t_{(16)}$ =3.05, p<0.05) and third (Mecp2- = 2.7 ± 0.2 , Mecp2+/+ = 3.7 ± 0.3 , Mecp2+ = 3.2 ± 0.2 ; $t_{(16)}$ =4.25, p<0.001) branch orders compared to both other genotypes (2-way repeated measures ANOVA and Bonferroni posttests, $F_{(16,96)}$ = 1.69, p=0.03). Mecp2- neurons therefore have, on average, one less branch point at each of the second and third branch orders. No statistically significant differences were found between Mecp2- and Mecp2+ cells at the first or fourth branch orders, yet Mecp2- cells appear to have fewer branch points here, raising the possibility that these branch orders may be affected by MECP2 mutation as well.

Partition asymmetry was next calculated at each branch order to assess whether branching patterns were altered in Mecp2- neurons (Figure 6C). No difference was found between the genotypes across branch orders (2-way repeated measures ANOVA and Bonferroni post-tests, $F_{(8,48)} = 2.23 \text{ p}=0.58$), indicating that Mecp2- neurons do not have more asymmetric branching patterns than Mecp2+ cells.

The number of branches per branch order was then calculated (Figure 6D) to confirm that the reduced numbers of second and third order branch points seen in Mecp2- neurons also results in a reduction in the number of subsequent branches in the third and fourth branch orders of these neurons. Mecp2- neurons have 16% fewer third order branches (~2 branches) (Mecp2- = 9.2 ± 0.7 , Mecp2+/+ = 11.0 ± 0.1 , Mecp2+ = 10.6 ± 0.7 ; $t_{(17)}$ =3.59, p<0.01) and 25% fewer fourth order branches (~2 branches) (Mecp2- = 5.3 ± 0.3 , Mecp2+/+ = 7.0 ± 0.7 , Mecp2+ = 6.2 ± 0.4 ; $t_{(14)}$ =3.46, p<0.01; 2-way repeated measures ANOVA and Bonferroni post-tests, $F_{(14,84)} = 1.51$ p=0.02). These data confirm that the lesser numbers of second and third order branch points result in a reduction in third and fourth order branches.

The total dendritic length per branch order was quantified (Figure 6E) to see if the reduced number of third and fourth order branches in Mecp2- neurons could account for the total reduction in dendritic length observed in these cells. Mecp2- neurons were found to have significantly less length in the third (17%) and fourth (27%) branch orders, as was predicted from the reduced number of third and fourth order branches (third order: Mecp2- = 499.8 ± 28.0 μ m, Mecp2+/+ = 599.5 ± 43.2 μ m, Mecp2+ = 554.6 ± 50.4 μ m; t₍₁₄₎=2.93, p<0.05; fourth order: Mecp2- = 300.3 ± 34.9 μ m, Mecp2+/+ = 414.0 ± 23.9 μ m, Mecp2+ = 380.2 ± 17.4 μ m; t₍₁₄₎=3.34, p<0.01; 2-way repeated measures ANOVA and Bonferroni post-tests, F_(14,84) = 1.15 p=0.03). These data indicate that Mecp2- cells have 100 μ m and 115 μ m less dendritic length in the third and fourth branch orders, respectively.

Finally, the average dendritic length per branch order was assessed (Figure 6F) to determine whether the shorter branches observed in Mecp2- neurons could also contribute to the reduced length observed in these cells. No difference was found in average branch length per branch order between the genotypes (2-way repeated measures ANOVA and Bonferroni post-tests, $F_{(8,48)} = 0.36 p=0.8$). This suggests that the shorter dendritic length observed in Mecp2- neurons is not caused by their having shorter branches than the other genotypes, but primarily from their reduced number of third and fourth order branches.

Cell autonomous effects on soma and nuclear areas

The nuclear and somatic areas of Mecp2+ and Mecp2- Layer V motor pyramidal neurons were measured in *Mecp2+/-* mice (Figure 7). The mean nuclear area of Mecp2- neurons in these experiments was 13% smaller than that of Mecp2+ neurons (n=24 animals, Mecp2- = 93.3 \pm 1.4 µm², Mecp2+ 107.9 \pm 1.2 µm², t₍₄₆₎=8.04, p<0.0001; Figure 9A). A plot of the entire population of individual nuclear areas (Mecp2+ n=598 neurons, Mecp2- n=418 neurons (Figure 7B) shows that the proportion of small neurons between the two genotypes are comparable, while the proportion of neurons in the largest size class is considerably smaller in the Mecp2- cell population than it is in the Mecp2+ population. The mean soma area of Mecp2- neurons is 15% smaller than Mecp2+ (n=24 animals, Mecp2- = 143.3 \pm 2.5 µm², Mecp2+ 169.2 \pm 1.9 µm², t₍₄₆₎=8.25, p<0.0001; Figure 7C). A graph of individual soma areas (Mecp2+ n=598 neurons, Mecp2- n=418 neurons, Figure 7D) shows that smallest somata of Mecp2- (79.8 µm²) and Mecp2+ (76.3 µm²) genotypes are comparable, while the largest quartile of MECP2- somata is considerably smaller than the largest quartile of Mecp2+ somata. Frequency distributions of the areas of nuclei (Figure 7E) and somata (Figure 7F) show that the distribution of Mecp2- neuron sizes is significantly shifted towards the smaller size classes compared to Mecp2+ neurons (nuclear area t_(3,21)=23.66, p<0.0001, soma area t_(3,36)=27.57, p<0.0001). To assess whether the soma or nuclear areas in

Mecp2- neurons were reduced disproportionally in relation to the each other, Mecp2+ and Mecp2- soma areas were plotted against their respective nuclear areas from within the same population (Figure 7G). The resulting slopes of Mecp2- and Mecp2+ populations were not significantly different ($F_{(1,1012)}$ =1.08, p=0.3) showing that while Mecp2- nuclei and soma are smaller in area than their Mecp2+ counterparts, the proportion of soma and nuclear areas are similar in both genotypes. *Symptom severity of Mecp2+/- mice across age and XCI ratios*

To determine the interaction between age, cell phenotype, and XCI ratios, animals were scored for their phenotype severity at their time of sacrifice for immunohistochemical staining (from 5-22 months of age). Within the cohort of mice selected for our study the phenotype severity was highly variable, with some mice dying prematurely as young as 2 months of age (i.e. before they could be used for experiments), and others surviving into old age (22 months) with only mild symptoms (Figure 8). Increasing age did not correlate with increased phenotype severity (p=0.2, $R^2=0.038$). To determine whether symptom levels vary according to XCI ratio, XCI ratios were analyzed by immunohistochemistry and compared against phenotype severity. It was found that XCI ratios were also not correlated with phenotype severity (Figure 8; p=0.66, $R^2=0.0062$). Surprisingly, mice with a highly skewed XCI ratio favouring expression of the wild-type chromosome (10% Mecp2-:90% Mecp2+) did not have less severe symptoms than mice with a more balanced XCI ratio (50% Mecp2-:50% Mecp2+). *Effect of age on soma and nuclear areas*

Age did not affect nuclear area (Mecp2+ $F_{(1,22)}=0.0089$, p=0.92, Mecp2- $F_{(1,22)}=0.012$, p=0.61) or somatic area (Mecp2+ $F_{(1,22)}=0.34$, p=0.57, Mecp2- $F_{(1,22)}=0.069$, p=0.79) for either genotype (Figure 9). The linear regression slopes for Mecp2+ and Mecp2- cells are not significantly different in either nuclear ($F_{(1,44)}=0.21$, p=0.65) or soma ($F_{(1,44)}=0.023$, p=0.88) areas, plotted across age from 5 to 22 month. *Effect of XCI ratios on soma and nuclear areas*

To determine if XCI ratios have an effect on soma or nuclear sizes of Mecp2+ and Mecp2neurons, somatic and nuclear areas were plotted against the proportion of Mecp2- neurons in the brain (Figure 10). The linear regression slope of Mecp2- nuclear area was significantly different from that of Mecp2+ nuclear area ($F_{(1,44)}$ =4.26, p=0.045; Figure 10A). This indicates that as the proportion of wildtype neurons in the brain increases, the nuclear area of Mecp2+ neurons increases, while the area of Mecp2- nuclei decreases.. The slope of the Mecp2- soma area does not differ statistically from that of the Mecp2+ soma area ($F_{(1,44)}$ =3.41, p=0.07; Figure 10B), however the trend of diverging slopes is apparent.

To minimize inter-animal differences (i.e. differences associated with tissue preparation and fixation), Mecp2- areas were normalized to Mecp2+ areas in each slice and plotted across the proportion of Mecp2- neurons in the brain (Figure 10C). The normalized nuclear area of Mecp2- neurons was positively correlated to the proportion of Mecp2- neurons in the brain ($F_{(1,22)}$ =18.14, p=0.0003). The

normalized soma area of Mecp2- neurons was also positively correlated to the proportion of Mecp2neurons in the brain ($F_{(1,22)}$ =8.08, p=0.0095). These data suggest that animals with a highly skewed XCI ratio, favouring expression of the wild-type allele (<30% Mecp2-/>70% Mecp2+) have the most severe Mecp2- neuronal phenotype (17-22% smaller than Mecp2+). Animals with a balanced XCI ratio (~50% Mecp2-/50% Mecp2+) have Mecp2- neurons with a comparatively less severe neuronal phenotype (11-17% smaller than Mecp2+).

Soma and nuclear size of Mecp2 +/+ versus Mecp2+ neurons

For the study summarized in figures 4-10, we did not have a comparison group of wild-type littermates, so it was not possible to test whether Mecp2+ neurons were comparable to Mecp2+/+ neurons, with respect to soma and nuclear size. Since a number of sources of error can potentially affect cell size, such as variation in cell shrinkage during fixation, we analyzed an entirely new group of mice comprising heterozygous females and their wild-type female littermates. As expected, comparison of Mecp2+ to Mecp2- soma and nuclear sizes in this new cohort of animals revealed a difference for the group mean data (Figure 11; mean soma difference 8.8 μ m², p < 0.0020, n=11; mean nuclear difference 4.1 μ m, p < 0.005). Pairing was effective (soma R²=0.97 p<0.0001, nuclear R²=0.90 p,0.0001), indicating that interanimal variability contributes a substantial amount of variance to the group means.

Mecp2+ neurons in the heterozygous brain were smaller, on average, than Mecp2+/+ neurons (mean soma difference 15.4 μ m², p < 0.032; nucleus 10.0 μ m², p< 0.038, n=10 WT, n= 11 Mecp2+). Pairwise comparisons based on littermate status (siblings) or day of tissue preparation did not measurably improve the statistical discrimination of the difference between Mecp2+/+ and Mecp2+ neurons. This suggests that the main source of variance in these experiments is intrinsic inter-animal variability in cell size, rather than variance in tissue preparation. Thus we found a sample size of more than 8 animals was needed to confidently avoid Type 2 errors when comparing between Mecp2+ neurons in *Mecp2^{+/+}* and *Mecp2^{+/+}* and *Mecp2^{+/+}* animals. When the overall cell size distribution is examined (Figure 11B), the tendency for the Mecp2- population to appear shifted toward smaller sizes when compared to the population of Mecp2+ cells also seems shifted toward small size when compared to the population of cells in *Mecp2^{+/+}* animals. Once again, the smallest cells in each population are similar in size.

DISCUSSION

MECP2- neurons have reduced total dendritic length and fewer third and fourth order branches

We find that Mecp2- layer V pyramidal neurons in heterozygous brains have shorter basal dendritic length, due to their having fewer third and fourth order branches compared to Mecp2+ neurons (see Figure 12 for summary). This supports previous research in *Mecp2*^{tm1.1Jae} male mice that also found a

reduction in the number of higher order branches in layer V cells (Stuss *et al.*, 2012). Within the basal dendritic compartment, the principal difference between neurons containing mutant MECP2 protein compared to those expressing the wild-type allele in both $Mecp2^{+/-}$ and $Mecp2^{+/+}$ animals is a selective reduction in the number of third and fourth order branches. The similar pattern of reduced basal dendritic morphology of Layer V pyramidal neurons for both Mecp2- neurons in $Mecp2^{+/-}$ females and male $Mecp2^{-/y}$ mice indicates that MECP2 acts predominantly in a cell-autonomous manner to affect the dendritic morphology of this compartment of Mecp2- neurons. These results are consistent with other studies that found that callosal projection neurons from $Mecp2^{-/y}$ mice transplanted into either wild-type or $Mecp2^{-/y}$ brains had primarily cell autonomous reductions in dendritic morphology (Kishi and Macklis, 2009).

Mecp2+ neurons were indistinguishable from neurons in $Mecp2^{+/+}$ animals with respect to cumulative dendritic length, number of branch points, and maximum branch order versus distance from the soma. Therefore, we find that the basic neuronal morphology of Layer V basal dendrites is primarily affected by MECP2 in a cell-autonomous manner. However, examination of some branch order analyses (Figure 6B, D, E) suggest Mecp2+ neurons have properties intermediate to those of Mecp2+/+ and Mecp2-. Therefore, while cell-autonomous (genotype-specific) effects appear to dominate branch structure, additional non-cell-autonomous effects may also be influencing branching. An inherent difficulty with analysis of basal dendrite branching in layer V cortical neurons is the large variation in dendritic structure inherent to this neuronal population (Tsiola *et al.*, 2003). A large number of neurons must therefore be sampled to obtain a representation of the "average" layer V pyramidal neuron making detailed comparisons between the dendritic arbours of neurons expressing wild-type MECP2 in $Mecp2^{+/+}$ and $Mecp2^{+/-}$ animals a challenging task.

Our analysis of 35 heterozygous mice did not reveal a correlation between XCI ratios in the primary motor cortex and the severity of behavioural phenotype, although previous research has suggested that the severity of the RTT phenotype may be influenced by the proportion of Mecp2- neurons in the brain (Sirianni *et al.*, 1998; Amir *et al.*, 2000; Plenge *et al.*, 2002; Miltenberger-Miltenyi and Laccone, 2003; Young and Zoghbi, 2004; Weaving *et al.*, 2005; Archer *et al.*, 2007). We acknowledge that our phenotype severity was scored using gross behavioural tests, rather than more extensive testing which has been shown to reveal subtle abnormalities even in young female mice (Samaco *et al.*, 2012), which do not ordinarily show obvious symptoms. Additionally, XCI ratios were determined using immunohistochemical analysis of Mecp2+ and Mecp2- neurons in the motor cortex of the brain, whereas other studies use blood or peripheral tissue to determine the XCI ratio, which has been reported to not necessarily correlate with the XCI in the brain (Gale *et al.*, 1994; Sharp *et al.*, 2000; Young and Zoghbi, 2004). Furthermore, although immunohistochemical determination of XCI may be more accurate, we

limited our analysis to layer V pyramidal neurons, which appear to be particularly subject to wide fluctuations in the pattern of XCI inactivation (Wu *et al.*, 2014).

Soma and nuclear size of Mecp2- neurons are reduced in the Mecp2+/- brain

Analysis of soma and nuclear size revealed that the average Mecp2- soma/nucleus is smaller than that of the average Mecp2+ neuron, consistent with previous studies comparing male $Mecp2^{-/y}$ to $Mecp2^{+/y}$ mice (Kishi and Macklis, 2004; Fukuda *et al.*, 2005). When the distribution of individual cell sizes from all animals are pooled, the Mecp2- population appears to primarily lack neurons comparable to the larger ones in the Mecp2+ population, while the prevalence of the smaller neurons in both genotypes is similar. This indicates that mutant MECP2 may limit the size of neurons, but does not decrease the size of Mecp2- somata beyond that of the smallest subset of Mecp2+ neurons.

Recent studies have found that MECP2 controls nuclear size in a largely cell-autonomous manner (Yazdani *et al.*, 2012; Singleton *et al.*, 2011). We find significant differences in the average sizes of somata and nuclei between Mecp2+ and Mecp2- neurons in the heterozygous brain, which appears to support this hypothesis. However, when comparing Mecp2+ cells in the heterozygous female brain to Mecp2+/+ cells in the wild-type brain, we also find a significant difference in cell size, with neurons from the wild-type brain being considerably larger. This suggests additional non-cell autonomous mechanisms for the regulation of cell size by MECP2, which may have a detrimental effect on Mecp2+ cell size in the heterozygous brain (Wu et al., 2014). This is an intriguing observation, since most studies to date have not examined neuron morphology in heterozygous females. Additionally, we found that while we needed only a small sample size to confidently avoid type 2 errors for comparisons between Mecp2+ and Mecp2-cells in heterozygous brains, a relatively large sample size (n>8) was required to demonstrate significant differences between Mecp2-positive cells in $Mecp2^{+/-}$ and $Mecp2^{+/+}$ littermate pairs. This may partially explain why we were able to note differences in Mecp2+ cell size between wildtype and heterozygous animals, but were not able to note similar non-cell autonomous differences in dendritic morphology. *An effect of XCI ratios on soma and nuclear areas*

Previous research has suggested that mutations in *Mecp2* may act non-cell-autonomously to affect dendritic morphology through the secretion of inhibitory substances by MECP2-deficient glia (Ballas *et al.*, 2009; Maezawa *et al.*, 2009; 2010). Braunschweig *et al.*, (2004) report that as the number of Mecp2neurons in the female $Mecp2^{+/-}$ brain increases, the expression of MECP2 in Mecp2+ neurons decreases proportionally, suggesting that in addition to cell-autonomous genotype-specific effects, *Mecp2* mutations can negatively impact neurons through actions of surrounding Mecp2- neurons and/or glia. When comparing Mecp2+/+ neuron size (soma and nuclear) in wild-type animals to that of Mecp2+ neurons in heterozygous females, (Figure 11) we see evidence of a non-cell-autonomous effect, in which the presence of Mecp2- cells in the brain is associated with reduced cell size for Mecp2+ neurons.

Comparing Mecp2+ to Mecp2- neurons across XCI ratios presents a more complex picture, however, particularly when between-animal differences are normalized by within-animal comparisons. As the proportion of Mecp2- neurons in the brain decreases Mecp2+ neurons tend to get larger (consistent with non-cell-autonomous effects). However Mecp2- soma and nuclear size tend to either not change or decrease, despite an increasing "wild-type" brain environment, resulting in a strong inverse relationship between XCI ratio and the ratio of Mecp2-/Mecp2+ cell size, shown in Figure 10C. Given that soma size has been reported to correlate with dendritic complexity (Kaufmann et al., 1995), these data may support a model in which Mecp2- neurons find it difficult to compete for connections in female brains with a highly skewed XCI ratio where the Mecp2+ neurons predominate. One hypothesis is that the consequent inability to compete for synaptic connections reduces the size of Mecp2- neurons to a greater degree than cell-autonomous effects alone. In contrast, in females with a balanced XCI ratio, the "Hebbian" synaptic environment is less competitive overall therefore allowing Mecp2- neurons to make more contacts with other neurons (Figure 13). Alternatively or additionally Mecp2- neurons may simply be unable to benefit from the more wild-type brain environment as the proportion of wild-type neurons increases with highly skewed XCI. Regardless of the cause, the result is that the cell-autonomous effects of MECP2 mutation dominate even in a predominantly wild-type environment.

Acknowledgments

The authors acknowledge the financial support of the International Rett Syndrome Foundation USA and the Natural Sciences and Engineering Research Council, Canada.

References cited

Abdala AP, Dutschmann M, Bissonnette JM and Paton JF (2010) Correction of respiratory disorders in a mouse model of Rett syndrome. Proc Natl Acad Sci U S A *107*, 18208-213.

Abràmoff MD, Magalhães PJ and Ram SJ (2004) Image processing with ImageJ. Biophotonics international 11, 36-42.

Adler DA, Quaderi NA, Brown SD, Chapman VM, Moore J, Tate P and Disteche CM (1995) The X-linked methylated DNA binding protein, Mecp2, is subject to X inactivation in the mouse. Mamm Genome *6*, 491-92.

Amir RE, Van den Veyver IB, Wan M, Tran CQ, Francke U and Zoghbi HY (1999) Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. Nat Genet 23, 185-88.

Amir RE, and Zoghbi HY (2000) Rett syndrome: methyl-CpG-binding protein 2 mutations and phenotype-genotype correlations. Am J Med Genet 97, 147-52.

Archer H, Evans J, Leonard H, Colvin L, Ravine D, Christodoulou J, Williamson S, Charman T, Bailey ME, Sampson J, de Klerk N and Clark A (2007) Correlation between clinical severity in patients with Rett syndrome with a p.R168X or p.T158M MECP2 mutation, and the direction and degree of skewing of X-chromosome inactivation. J Med Genet *44*, 148-52.

Armstrong D, Dunn JK, Antalffy B and Trivedi R (1995) Selective dendritic alterations in the cortex of Rett syndrome. J Neuropathol Exp Neurol 54, 195-201.

Armstrong DD (1995). The neuropathology of Rett syndrome--overview 1994. Neuropediatrics 26, 100-04.

Ballas N, Lioy DT, Grunseich C and Mandel G. (2009) Non-cell autonomous influence of Mecp2deficient glia on neuronal dendritic morphology. Nat Neurosci 12, 311-17.

Ballesteros-Yáñez, I., Benavides-Piccione R, Elston GN, Yuste R and DeFelipe J (2006) Density and morphology of dendritic spines in mouse neocortex. Neuroscience 138, 403-09.

Balmer D, Goldstine J, Rao YM and LaSalle JM (2003) Elevated methyl-CpG-binding protein 2 expression is acquired during postnatal human brain development and is correlated with alternative polyadenylation. J Mol Med *81*, 61-68.

Bauman ML, Kemper TL and Arin DM (1995). Pervasive neuroanatomic abnormalities of the brain in three cases of Rett's syndrome. Neurology 45, 1581-86.

Belichenko PV, Oldfors A, Hagberg B and Dahlström A (1994) Rett syndrome: 3-D confocal microscopy of cortical pyramidal dendrites and afferents. Neuroreport *5*, 1509-513.

Belichenko PV, Hagberg B and Dahlström A (1997) Morphological study of neocortical areas in Rett syndrome. Acta Neuropathol 93, 50-61.

Belichenko NP, Belichenko PV and Mobley WC (2009) Evidence for both neuronal cell autonomous and nonautonomous effects of methyl-CpG-binding protein 2 in the cerebral cortex of female mice with Mecp2 mutation. Neurobiol Dis *34*, 71-77.

Belmont JW (1996) Genetic control of X inactivation and processes leading to X-inactivation skewing. Am J Hum Genet 58, 1101-08.

Bienvenu T and Chelly J (2006) Molecular genetics of Rett syndrome: when DNA methylation goes unrecognized. Nat Rev Genet 7, 415-26.

Bissonnette JM and Knopp SJ (2006) Separate respiratory phenotypes in methyl-CpG-binding protein 2 (Mecp2) deficient mice. Pediatr Res 59, 513-18.

Bissonnette JM, Knopp SJ, Maylie J, and Thong T (2007) Autonomic cardiovascular control in methyl-CpG-binding protein 2 (Mecp2) deficient mice. Auton Neurosci 136, 82-89.

Braunschweig D, Simcox T, Samaco RC and LaSalle JM (2004) X-Chromosome inactivation ratios affect wild-type Mecp2 expression within mosaic Rett syndrome and Mecp2-/+ mouse brain. Hum Mol Genet *13*, 1275-286.

Brown CJ and Robinson WP (2000). The causes and consequences of random and non-random X chromosome inactivation in humans. Clin Genet *58*, 353-363.

Carter JC, Lanham DC, Pham D, Bibat G, Naidu S and Kaufmann WE (2008) Selective cerebral volume reduction in Rett syndrome: a multiple-approach MR imaging study. AJNR Am J Neuroradiol 29, 436-41.

Chandler SP, Guschin D, Landsberger N and Wolffe AP (1999) The methyl-CpG binding transcriptional repressor Mecp2 stably associates with nucleosomal DNA. Biochemistry *38*, 7008-018.

Chapleau CA, Calfa GD, Lane MC, Albertson AJ, Larimore JL, Kudo S, Armstrong DL, Percy AK and Pozzo-Miller L (2009). Dendritic spine pathologies in hippocampal pyramidal neurons from Rett syndrome brain and after expression of Rett-associated MECP2 mutations. Neurobiol Dis *35*, 219-33.

Charman T, Neilson TC, Mash V, Archer H, Gardiner MT, Knudsen GP, McDonnell A, Perry J, Whatley SD, Bunyan DJ, Ravn K, Mount RH, Hastings RP, Hulten M, Orstavik KH, Reilly S, Cass H, Clarke A, Kerr AM, Bailey ME (2005). Dimensional phenotypic analysis and functional categorisation of mutations reveal novel genotype-phenotype associations in Rett syndrome. Eur J Hum Genet *13*, 1121-130.

n

Chen RZ, Akbarian S, Tudor M and Jaenisch R (2001) Deficiency of methyl-CpG binding protein-2 in CNS neurons results in a Rett-like phenotype in mice. Nat Genet 27, 327-31.

Costa LDF, Zawadzki K, Miazaki M, Viana MP and Taraskin SN (2010). Unveiling the neuromorphological space. Front Comput Neurosci 4, 150.

Cuddapah VA, Pillai RB, Shekar KV, Lane JB, Motil KJ, Skinner SA, Tarquinio DC, Glaze DG, McGwin G, Kaufmann WE, Percy AK, Neul JL, Olsen ML (2014). Methyl-CpG-Binding Protein 2 (MECP2) mutation type is associated with disease severity in Rett Syndrome. J Med Genet *51 (3)*: 152-58.

Derecki NC, Cronk JC, Lu Z, Xu E, Abbott SB, Guyenet PG and Kipnis J (2012) Wild-type microglia arrest pathology in a mouse model of Rett syndrome. Nature 484, 105-09.

Feng G, Mellor RH, Bernstein M, Keller-Peck C, Nguyen QT, Wallace M, Nerbonne JM, Lichtman JW and Sanes JR (2000) Imaging neuronal subsets in transgenic mice expressing multiple spectral variants of GFP. Neuron 28, 41-51.

Fukuda T, Itoh M, Ichikawa T, Washiyama K and Goto Y (2005) Delayed maturation of neuronal architecture and synaptogenesis in cerebral cortex of Mecp2-deficient mice. J Neuropathol Exp Neurol *64*, 537-544.

Gale RE, Wheadon H, Boulos P and Linch DC (1994) Tissue specificity of X-chromosome inactivation patterns. Blood *83*, 2899-905.

Ghosh RP, Horowitz-Scherer RA, Nikitina T, Shlyakhtenko LS and Woodcock CL (2010) Mecp2 binds cooperatively to its substrate and competes with histone H1 for chromatin binding sites. Mol Cell Biol *30*, 4656-670.

Guy J, Hendrich B, Holmes M, Martin JE and Bird A (2001) A mouse Mecp2-null mutation causes neurological symptoms that mimic Rett syndrome. Nat Genet *27*, 322-26.

Guy J, Gan J, Selfridge J, Cobb S and Bird A (2007) Reversal of neurological defects in a mouse model of Rett syndrome. Science *315*, 1143-47.

Guy J, Cheval H, Selfridge J and Bird A (2010) The Role of Mecp2 in the Brain. Annu Rev Cell Dev Biol 27, 631-52.

Jordan C, Li HH, Kwan HC and Francke U (2007) Cerebellar gene expression profiles of mouse models for Rett syndrome reveal novel Mecp2 targets. BMC Med Genet *8*, 36.

Kaufmann, W.E., Naidu, S., and Budden, S. (1995). Abnormal expression of microtubule-associated protein 2 (MAP-2) in neocortex in Rett syndrome. Neuropediatrics *26*, 109-113.

Kaufmann WE, MacDonald SM and Altamura CR (2000). Dendritic cytoskeletal protein expression in mental retardation: an immunohistochemical study of the neocortex in Rett syndrome. Cereb Cortex *10*, 992-1004.

Kaufmann WE, Tierney E, Rohde CA, Suarez-Pedraza MC, Clarke MA, Salorio CF, Bibat G, Bukelis I, Naram D, Lanham DC and Naidu S (2011) Social impairments in Rett syndrome: characteristics and relationship with clinical severity. J Intellect Disabil Res *56*, 233-47.

Kishi N and Macklis JD (2004) MECP2 is progressively expressed in post-migratory neurons and is involved in neuronal maturation rather than cell fate decisions. Mol Cell Neurosci 27, 306-21.

Kishi N and Macklis JD (2009) Mecp2 functions largely cell-autonomously, but also non-cellautonomously, in neuronal maturation and dendritic arborization of cortical pyramidal neurons. Exp Neurol 222, 51-58.

Kriaucionis S, Paterson A, Curtis J, Guy J, Macleod N and Bird A (2006). Gene expression analysis exposes mitochondrial abnormalities in a mouse model of Rett syndrome. Mol Cell Biol 26, 5033-042.

Laurvick CL, de Klerk N, Bower C, Christodoulou J, Ravine D, Ellaway C, Williamson S and Leonard H (2006) Rett syndrome in Australia: a review of the epidemiology. J Pediatr *148*, 347-52.

Lioy DT, Garg SK, Monaghan CE, Raber J, Foust KD, Kaspar BK, Hirrlinger PG, Kirchhoff F, Bissonnette JM, Dallas N and Mandel G (2011) A role for glia in the progression of Rett's syndrome. Nature 475, 497-500.

Longair MH, Baker DA and Armstrong JD (2011) Simple Neurite Tracer: open source software for reconstruction, visualization and analysis of neuronal processes. Bioinformatics 27, 2453-54.

Maezawa I and Jin LW (2010) Rett syndrome microglia damage dendrites and synapses by the elevated release of glutamate. J Neurosci *30*, 5346-356.

Maezawa I, Swanberg S, Harvey D, LaSalle JM and Jin LW (2009) Rett syndrome astrocytes are abnormal and spread Mecp2 deficiency through gap junctions. J Neurosci 29, 5051-061.

Metcalf BM, Mullaney BC, Johnston MV and Blue ME (2006) Temporal shift in methyl-CpG binding protein 2 expression in a mouse model of Rett syndrome. Neuroscience 139, 1449-460.

Miltenberger-Miltenyi G and Laccone F (2003) Mutations and polymorphisms in the human methyl CpG-binding protein MECP2. Hum Mutat 22, 107-15.

Nan X, Campoy FJ and Bird A.(1997) Mecp2 is a transcriptional repressor with abundant binding sites in genomic chromatin. Cell *88*, 471-81.

Neul JL, Kaufmann WE, Glaze DG, Christodoulou J, Clarke AJ, Bahi-Buisson N, Leonard H, Bailey ME, Schanen NC, Zappella M, Reineri A, Huppke P and Percy AK (2010) Rett syndrome: Revised diagnostic criteria and nomenclature. Ann Neurol *68*, 944-50.

Nikitina T, Ghosh RP, Horowitz-Scherer RA, Hansen JC, Grigoryev SA and Woodcock CL (2007) Mecp2-chromatin interactions include the formation of chromatosome-like structures and are altered in mutations causing Rett syndrome. J Biol Chem 282, 28237-245.

Nuber UA, Kriaucionis S, Roloff TC, Guy J, Selfridge J, Steinhoff C, Schulz R, Lipkowitz B, Ropers HH, Homes Mc and Bird A (2005) Up-regulation of glucocorticoid-regulated genes in a mouse model of Rett syndrome. Hum Mol Genet *14*, 2247-256.

Philippe C, Villard L, De Roux N, Raynaud M, Bonnefond JP, Pasquier L, Lesca G, Mancini J, Jonveaux P, Moncini J, Jonveaux P, Moncla A, ChellJ and Bienvenu T (2006) Spectrum and distribution of MECP2 mutations in 424 Rett syndrome patients: a molecular update. Eur J Med Genet *49*, 9-18.

Plenge RM, Stevenson RA, Lubs HA, Schwartz CE and Willard HF (2002) Skewed X-chromosome inactivation is a common feature of X-linked mental retardation disorders. Am J Hum Genet 71, 168-73.

Puck JM and Willard HF (1998) X inactivation in females with X-linked disease. N Engl J Med 338, 325-28.

Reiss AL, Faruque F, Naidu S, Abrams M, Beaty T, Bryan RN and Moser H (1993). Neuroanatomy of Rett syndrome: a volumetric imaging study. Ann Neurol *34*, 227-34.

Rett A (1966) [On a unusual brain atrophy syndrome in hyperammonemia in childhood]. Wien Med Wochenschr 116, 723-26.

Rodriguez A, Ehlenberger D, Kelliher K, Einstein M, Henderson SC, Morrison JH, Hof PR and Wearne SL (2003) Automated reconstruction of three-dimensional neuronal morphology from laser scanning microscopy images. Methods *30*, 94-105.

Ross MT, Grafham DV, Coffey AJ, Scherer S, McLay K, Muzny D, Platzer M, Howell GR, Burrows C, *et al.*, (2005). The DNA sequence of the human X chromosome. Nature *434*, 325-37.

Samaco RC, Mandel-Brehm C, Chao HT, Ward CS, Fyffe-Maricich SL, Ren J, Hyland K, Thaller C, Maricich SM, Humphreys P, Greer JJ, Percy A, Glaze DG, Zoghbi HY and Neul JL (2009) Loss of Mecp2 in aminergic neurons causes cell-autonomous defects in neurotransmitter synthesis and specific behavioral abnormalities. Proc Natl Acad Sci USA *106*, 21966-971.

Samaco RC, McGraw CM, Ward CS, Sun Y, Neul JL and Zoghbi HY (2012). Female *Mecp2+/-* mice display robust behavioral deficits on two different genetic backgrounds providing a framework for preclinical studies. Hum Mol Genet. 22(1), 96-109

Santos M, Silva-Fernandes A, Oliveira P, Sousa N and Maciel P (2007) Evidence for abnormal early development in a mouse model of Rett syndrome. Genes Brain Behav *6*, 277-286.

Schnell SA, Staines WA and Wessendorf MW (1999) Reduction of Lipofuscin-like autofluorescence in fluorescently labeled tissue. Journal of Histochem & Cytochem 47(8), 719-3-.

Scorcioni R, Polavaram S and Ascoli GA (2008) L-Measure: a web-accessible tool for the analysis, comparison and search of digital reconstructions of neuronal morphologies. Nat Protoc *3*, 866-76.

Shahbazian MD, Antalffy B, Armstrong DL and Zoghbi HY (2002) Insight into Rett syndrome: Mecp2 levels display tissue- and cell-specific differences and correlate with neuronal maturation. Hum Mol Genet 11, 115-24.

Sharp A, Robinson D and Jacobs P (2000) Age-and tissue-specific variation of X chromosome inactivation ratios in normal women. Hum Genet *107*, 343-49.

Sholl DA (1953) Dendritic organization in the neurons of the visual and motor cortices of the cat. J Anat 87, 387-405.

Singleton MK, Gonzales ML, Leung KN, Yasui DH, Schroeder DI, Dunaway K and LaSalle JM (2011) Mecp2 is required for global heterochromatic and nucleolar changes during activity-dependent neuronal maturation. Neurobiol Dis *43*, 190-200.

Sirianni N, Naidu S, Pereira J, Pillotto RF and Hoffman EP (1998) Rett syndrome: confirmation of X-linked dominant inheritance, and localization of the gene to Xq28. Am J Hum Genet *63*, 1552-58.

Skene PJ, Illingworth RS, Webb S, Kerr AR, James KD, Turner DJ, Andrews R and Bird AP (2010) Neuronal Mecp2 is expressed at near histone-octamer levels and globally alters the chromatin state. Mol Cell *37*, 457-68.

Smrt RD, Pfeiffer RL and Zhao X (2011) Age-dependent expression of Mecp2 in a heterozygous mosaic mouse model. Hum Mol Genet 20, 1834-843.

Stuss DP, Boyd JD, Levin DB and Delaney KR (2012) Mecp2 Mutation Results in Compartment-Specific reductions in dendritic branching and spine density in Layer 5 Motor cortical neurons of YFP-H Mice. PLoS One 7, e31896.

Stuss DP, Manjinder C, Ng MK, de Paz AM, Williamson B, Missiaen K, Cosman JD, McPhee D, Esteller M, Hendel M, Delaney K and Ausio J (2013) Impaired *in vivo* binding of Mecp2 to chromatin in the absence of its DNA methyl-binding domain. *Nuc Acid Res.* 41(9) 4888-900

Subramaniam B, Naidu S and Reiss AL (1997) Neuroanatomy in Rett syndrome: cerebral cortex and posterior fossa. Neurology 48, 399-407.

Tsiola A, Hamzei-Sichani F, Pertlin Z and Yuste R (2003) Quantitative morphologic classification of layer 5 neurons from mouse primary visual cortex. J Comp Neurol *461*: 415-28

Wearne SL, Rodriguez A, Ehlenberger DB, Rocher AB, Henderson SC and Hof PR (2005) New techniques for imaging, digitization and analysis of three-dimensional neural morphology on multiple scales. Neuroscience *136*, 661-80.

Weaving LS, Christodoulou J, Williamson SL, Friend KL, McKenzie OL, Archer H, Evans J, Clarke A, Pelka GJ, Tam PP, Watson C, Lahooti H, Ellaway CJ, Bennetts B, Leonard H and Gécz J (2004) Mutations of CDKL5 cause a severe neurodevelopmental disorder with infantile spasms and mental retardation. Am J Hum Genet *75*, 1079-093.

Weaving LS, Ellaway CJ, Gécz J and Christodoulou, J (2005) Rett syndrome: clinical review and genetic update. J Med Genet 42, 1-7.

Wu H, Luo J, Yu H, Rattner A, Mo A, Wang Y, Smallwood PM, Erlanger B, Wheelan SJ, and Nathans J (2014) Cellular Resolution Maps of X Chromosome Inactivation: Implications for Neural Development, Function and Disease. Neuron *81*, 103-19.

Yazdani M, Deogracias R, Guy J, Poot RA, Bird A and Barde YA (2012) Disease Modeling Using Embryonic Stem Cells: Mecp2 Regulates Nuclear Size and RNA Synthesis in Neurons. Stem Cells *30*, 2128-139.

Young JI and Zoghbi HY (2004) X-chromosome inactivation patterns are unbalanced and affect the phenotypic outcome in a mouse model of rett syndrome. Am J Hum Genet 74, 511-20.

Zappella M, Meloni I, Longo I, Hayek G and Renieri A (2001) Preserved speech variants of the Rett syndrome: molecular and clinical analysis. Am J Med Genet *104*, 14-22.

Zappella M, Meloni I, Longo I, Canitano R, Hayek G, Rosaia L, Mari, F and Renieri A (2003) Study of MECP2 gene in Rett syndrome variants and autistic girls. Am J Med Genet B Neuropsychiatr Genet *119B*, 102-07.

Figure 1. Single cells microinjected with Alexa Fluor 594 dye.

Neurons selected for filling were distributed across brain slices to maximize the total number of visible cells per slice while preventing overlap of dendrites. With excitation and emission applied for AF594 imaging, YFP-expressing neurons were faintly visible, confirming that filled neurons were located in Layer V. Scale bar = 100µm.

Figure 2. Immunohistochemical genotyping to identify dye-filled, nuclear- or cytoplasm-counterstained neurons for morphological comparisons.

(A) C-terminal Mecp2 antibody staining appears punctate in Mecp2+ neurons and diffuse in Mecp2- neurons (arrows). (B) N-terminal Mecp2 staining appears punctate in Mecp2+ neurons, and is absent in Mecp2- neurons (arrows). (C) Alexa Fluor 594 (AF594) iontophoretically injected into lightly-fixed neurons fills the cell body and dendrites. (D) The composite of all channels reveals the genotype of the desired neuron (AF594 is shown in green pseudocolour for illustration purposes, pink = Mecp2+, red = Mecp2-). Scale bar = 20 μ m. E) DAPI nuclear labelling reveals which cells are in the in focal plane and delineates the boundary of nucleus. (F) Punctate (arrowhead) or diffuse (arrow) staining of the C-terminal MECP2 antibody differentiates Mecp2+ and Mecp2- cells, respectively. (G) The cytoplasm of neurons is labeled with Neurotrace Nissl body stain. (H) A composite image reveals the genotype of each neuron. Mecp2- cells appear blue/green because the red C-terminal MECP2 stain is faint and diffuse. Scale bar = 20 μ m. (I) Granular lipofuscin autofluorescence is faint in a brain slice at 9 months of age. (J) Lipofuscin accumulates in brain tissue over time. In an 18-month old animal strong autofluorescence makesthe visualization and quantification of fluorescent dyes difficult. (K) Treatment of slices with 5mM CuSO₄ in 50 mM ammonium acetate considerably reduces lipofuscin autofluorescence without affecting AF594 staining. Scale bar = 50 μ m.

Figure 3. Diagram of morphometric measurements.

A *primary dendrite* is any dendrite originating from the cell body. A *branch point* occurs when the original dendrite bifurcates into two or more daughter trees. A centrifugal labelling scheme was used for assigning *branch order*, where the lowest order is given to the primary dendrites and increases when a branch point is reached. *Maximum branch order* is the maximum value reached by any primary tree on a given cell. A *branch* is a length of dendrite between the soma and a branch point, between two branch points, or between a branch point and a tip. A *tip* is the termination of any branch. *Sholl radii* are concentric spheres originating from the centroid of the soma and increasing at 10 µm intervals. *Maximum radial distance* is the maximum Sholl radius reached by any single dendrite per cell. *Partition asymmetry* is calculated at each branch point as $|T_1-T_2| / (T_1+T_2-2)$ where T_1 and T_2 are the number of tips of the two daughter trees, with values of one and zero indicating asymmetrical and symmetrical partitioning respectively. *Contraction* is calculated for each branch as a ratio between the Euclidean distance and the distance along the branch path. A value of one indicates perfectly straight dendrites, while lower values indicate increasing levels of tortuosity or winding of the dendrites. Derived in part from (Costa *et al.*, 2010).

Figure 4. Total dendritic length of MECP2- neurons is reduced

(A) Mecp- neurons have decreased dendritic length at distances of 70 to 130 μ m from the soma compared to both Mecp2+/+ and Mecp2+ neurons. The region between the dashed lines indicates where the slopes of Mecp2- and Mecp2+ cells deviate. (B) Mecp2- neuron total basal dendritic length is 15% less (275 μ m) compared to both Mecp2+/+ and Mecp2+ neurons. Dashed lines indicate the average maximum radial distance reached for each genotype. No significant difference was found between the genotypes in this measure. *p<0.05, **p<0.01, and ***p<0.001.

Figure 5. Mecp2- neurons have a reduction in the number of total branch points and maximum branch order across Sholl radii.

(A) Basal dendrites of all genotypes have similar path distances across Sholl radii. (B) All genotypes have the same number of primary dendrites. (C) Mecp2- neurons have an average of 3 fewer total branch points (a 17% reduction) compared to both other genotypes. The region between the dashed lines indicates where the Mecp2- neurons differ from Mecp2+ neurons in trend. (D) The maximum branch order reached by Mecp2- neurons at distances of 60 and 110 μ m from the soma is lower than both other genotypes. *p<0.05 and **p<0.01.

Figure 6. MECP2- neurons have fewer 3rd and 4th order branches compared to both other genotypes. (A)

Branch order diagram depicting the increasing order values with each successive branch point from the cell body. Symmetrical and asymmetrical branching patterns are shown (calculated at the second order branch points) which result in partition asymmetry values of 0 and 1, respectively. (**B**) Mecp2- neurons have fewer second and third order branch points compared to both genotypes. (**C**) Branching patterns in Mecp2- neurons are unaltered. (**D**) Mecp2- neurons have fewer third and fourth order branches. (**E**) Mecp2- neurons have reduced dendritic length in the third and fourth branch orders. (**F**) The average length of Mecp2- branches per branch order does not differ from that of Mecp2+ branches. *p<0.05, **p<0.01 and p<0.001.

Figure 7. Mecp2- neurons have smaller somata and nuclei than Mecp2+ neurons.

Mecp2- nuclei (A) and somata (C) are on average 13% and 15% smaller than Mecp2+ nuclei and somata, respectively (n=24 animals). The population of Mecp2- nuclei (B) and somata (D) have fewer large nuclei and somata compared to Mecp2+ (Mecp2+ n=598 cells and Mecp2- n=418 cells). Frequency distributions of nuclear (E) and soma (F) areas are significantly different between the genotypes (p<0.0001, nuclear area Mecp2+ R^2 =0.96 and Mecp2- R^2 =0.98. Soma area Mecp2+ R^2 =0.93 and Mecp2- R^2 =0.94). (G) Soma area is a predictor of nuclear area in both genotypes. Slopes are equal between the genotypes (p=0.30, Mecp2+ R^2 =0.79 and Mecp2- R^2 =0.78). ***p<0.001.

Figure 8. Age, phenotype severity and XCI ratios are not correlated in *Mecp2+/-* mice.

Increasing age is not a predictor of phenotype severity of *Mecp2+/-* mice. Aged *Mecp2+/-*animals are not more likely to have highly skewed XCI ratios (10-30% mutant neurons, red circles) and XCI ratios do not predict phenotype severity at any age. Premature death (x's) occurred across both young and old age groups. A premature death is defined as any animal that died spontaneously (not sacrificed for experiments), therefore XCI ratios were unavailable.

Figure 9. Age does not affect nuclear and soma areas.

(A) Nuclear area in Mecp2+ and Mecp2- neurons is not altered across age (Mecp2+ R^2 = 0.00040 and Mecp2- R^2 =0.012). (B) Soma area in both genotypes is not affected by age (Mecp2+ R^2 = 0.015 and Mecp2- R^2 =0.0031)

Figure 10. Nuclear and soma areas are correlated with XCI ratios.

(A) Nuclear areas of Mecp2+ and Mecp2- neurons diverge as XCI ratios become more skewed in favour of Mecp2+ expression (right to left). Slopes are significantly different between the genotypes (p=0.04, Mecp2+ R^2 =0.069 and Mecp2- R^2 =0.11). (B) Soma areas of Mecp2+ and Mecp2- neurons appear to diverge as the proportion of Mecp2- neurons in the brain decreases (right to left). Slopes are not significantly different between the two genotypes (p=0.07, Mecp2+ R^2 =0.13 and Mecp2- R^2 =0.041). (C) Mecp2- soma and nuclear areas were normalized to paired Mecp2+ values from the same animal. Mecp2- nuclei and somata in brains with XCI ratios highly skewed towards wild-type (>70%% Mecp2+) are significantly smaller than Mecp2- nuclei and somata in brains with more balanced XCI ratios (~50% Mecp2+). Slopes are significantly non-zero (nuclear area p=0.0003 R^2 =0.45, soma area p=0.0095 R^2 =0.29).

Figure 11. Soma and nuclear size comparison between wild-type and heterozygous females. (A & C)

Mecp2+/+ Layer V neurons (wildtype) are larger than Mecp2+ neurons in heterozygous mice, which suggests a non-cell-autonomous effect of Mecp2 deficiency (Mecp2+/+, n=10; Mecp2+, n=11; Mecp2- n=11). (**B & D**) Mecp2+ and Mecp2- neuronal populations in heterozygous brains each have fewer large cells than the population of Mecp2+/+ (wildtype) brains. The Mecp2- cell population also has fewer large cells than the Mecp2+ population in heterozygous brains. (Mecp2+/+, n=761 neurons; Mecp2+, n=482; Mecp2- n=502)

Figure 12. Diagram of average $Mecp2^{+/+}$, Mecp2+, and Mecp2- layer V pyramidal neurons. The average Mecp2+/+ and Mecp2+ neurons are equivalent across all morphological parameters investigated except soma and nuclear size. The average Mecp2- neuron (right) has approximately 15% less total dendritic length compared to either of the other genotypes. The average Mecp2- neuron has three fewer branch points, leading to a reduced number of higher order branches. The maximum branch order of Mecp2- neurons is also reduced. Mecp2- neurons have the same average maximum radial distance as both Mecp2+/+ and Mecp2+ neurons. Branch order: first = red, second = orange, third = green, fourth = blue, fifth = purple, sixth = black. Radii are 20 μ m.

Figure 13. Proposed model for cell non-autonomous effects of balanced and skewed XCI ratios. (A) Cell-

autonomous effects on the morphology of Mecp2- containing neurons, including reductions in dendritic length and fewer branch points compared to Mecp2+ cells, may render them less competitive for synaptic connections in the heterozygous brain. **(B)** A balanced XCI ratio may provide Mecp2- neurons more opportunity to make connections, as there are fewer large Mecp2+ neurons to compete with. This may reduce the severity of the morphological phenotype of Mecp2- cells. **(C)** A skewed XCI ratio favouring Mecp2+ neurons provides for a more competitive environment for synaptic connections. Mecp2- neurons are less able to compete and therefore have a more severe morphological phenotype.



















Phenotype Severity









